

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵: A61K 35/14, 37/00, 39/00, C07K 3/00, 13/00, 15/00, 17/00, C12P 21/06	A1	(11) International Publication Number: WO 94/28912 (43) International Publication Date: 22 December 1994 (22.12.94)
(21) International Application Number: PCT/US94/06701 (22) International Filing Date: 10 June 1994 (10.06.94) (30) Priority Data: 08/076,071 10 June 1993 (10.06.93) US (71) Applicants: THE REGENTS OF THE UNIVERSITY OF MICHIGAN [US/US]; 3003 S. State Street, Wolverine Tower, Room 2071, Ann Arbor, MI 48109-1280 (US). THE UNITED STATES OF AMERICA, represented by THE SECRETARY OF NAVY [US/US]; Naval Medical Research and Development Command, Building 1, Tower 12, 8901 Wisconsin Avenue, Bethesda, MD 20889-5606 (US). (72) Inventors: THOMPSON, Craig, B.; 12 Ridgeway, Ann Arbor, MI 48104 (US). JUNE, Carl, H.; 7 Harlow Court, Rockville, MD 20850 (US). (74) Agents: KENNEDY, Bill et al.; Morrison & Foerster, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: CD28 PATHWAY IMMUNOSUPPRESSION (57) Abstract The method of immunotherapy of the present invention involves the regulation of the T cell immune response through the activation or suppression/inactivation of the CD28 pathway. Induction of activated T cell lymphokine production occurs upon stimulatory binding of the CD28 surface receptor molecule, even in the presence of conventional immunosuppressants. Inhibition of CD28 receptor binding to an appropriate stimulatory ligand or inactivation of the CD28 signal transduction pathway through other means down-regulates CD28-pathway related T cell lymphokine production and its resulting effects.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

CD28 PATHWAY IMMUNOSUPPRESSION RELATED APPLICATIONS

This is a continuation-in-part of International Application Serial No. PCT/US93/03155, entitled "CD28 Pathway Immunoregulation," filed April 6, 1993 by Thompson *et al.*, which is a continuation of U.S. Application Serial No. 07/864,805, entitled "CD28 Pathway Immunoregulation," U.S. Application Serial No. 07/864,807, entitled "Immunotherapy Involving Stimulation of T_HCD28 Lymphokine Production," and U.S. Application Serial 07/864,866, entitled "Enhancement of CD28-Related Immune Response," all filed April 7, 1992 by Thompson, *et al.*, which are continuations-in-part of U.S. Serial No. 07/275,433, entitled "Immunotherapy Involving CD28 Stimulation," filed November 23, 1988 by Thompson *et al.*, now abandoned, and is also a continuation-in-part of International Application Serial No. PCT/US89/05304 (Publication No. WO 90/05541), entitled "Immunotherapy Involving CD28 Stimulation," filed November 22, 1989 by Thompson *et al.* and U.S. Patent No. 07/902,467 entitled "Immunotherapy Involving CD28 Stimulation," filed June 19, 1992 by Thompson *et al.*, all herein incorporated by reference.

SPONSORSHIP

Work on this invention was supported in part by Naval Medical Research and Development Command, Research Task No. M0095.0003-1007. The Government has certain rights in the invention.

BIOLOGICAL DEPOSIT

Murine hybridoma cell line 9.3 has been deposited with the American Type Culture Collection in Rockville, MD, in compliance with the provisions of the Budapest Treaty, and has been assigned ATCC Designation No. HB10271.

FIELD OF THE INVENTION

The present invention generally relates to immunotherapy. More particularly, the present invention relates to immunotherapy involving regulation of the CD28 T cell surface molecule.

BACKGROUND OF THE INVENTION

Thymus derived lymphocytes, referred to as T cells, are important regulators of *in vivo* immune responses. T cells are involved in cytotoxicity and delayed type hypersensitivity (DTH), and provide helper functions for B lymphocyte antibody production. In addition, T cells produce a variety of

- 2 -

lymphokines which function as immunomodulatory molecules, such as for example, interleukin-2 (IL-2), which can facilitate the cell cycle progression of T cells; tumor necrosis factor- α (TNF- α) and lymphotoxin (LT), cytokines shown to be involved in the lysis of tumor cells; Interferon- γ (IFN- γ), which displays a wide
5 variety of anti-viral and anti-tumor effects; and IL-3 and granulocyte-macrophage colony stimulating factor (GM-CSF), which function as multilineage hematopoietic factors.

Current immunotherapeutic treatments for diseases such as cancer, acquired immunodeficiency syndrome (AIDS) and attending infections, involve
10 the systemic administration of lymphokines, such as IL-2 and IFN- γ , in an attempt to enhance the immune response. However, such treatment results in non-specific augmentation of the T cell-mediated immune response, since the lymphokines administered are not specifically directed against activated T cells proximate to the site of infection or the tumor. In addition, systemic infusions
15 of these molecules in pharmacologic doses leads to significant toxicity. Present therapies for immunodeficient or immunodepressed patients also involve non-specific augmentation of the immune system using concentrated γ -globulin preparations. The stimulation of the *in vivo* secretion of immunomodulatory factors has not, until now, been considered a feasible alternative due to the
20 failure to appreciate the effects and/or mechanism and attending benefits of such therapy.

It would thus be desirable to provide a method of immunotherapy which enhances the T cell-mediated immune response and which is directed specifically toward T cells activated by an antigen produced by the targeted cell. It would
25 further be desirable to provide a method of immunotherapy which could take advantage of the patient's natural immunospecificity. It would also be desirable to provide a method of immunotherapy which can be used in immunodepressed patients. It would additionally be desirable to provide a method of immunotherapy which does not primarily rely on the administration of
30 immunomodulatory molecules in amounts having significant toxic effects.

It would also be desirable to provide a method of immunotherapy which, if so desired, could be administered directly without removal and reintroduction of T cell populations. It would further be desirable to provide a method of immunotherapy which could be used not only to enhance, but to suppress T

- 3 -

cell-mediated immunoresponses where such immun suppression would be advantage us, for example, in transplant patients, in patients exhibiting shock syndrome and in certain forms of autoimmune disease.

SUMMARY OF THE INVENTION

5 The present invention comprises a method of immunotherapy in which the T cell-mediated immune response is regulated by the CD28 pathway. Binding of the CD28 receptor with anti-CD28 antibodies or other stimulatory binding equivalents induces activated T cell-mediated lymphokine production. Immunosuppression or down-regulation is achieved by preventing CD28 receptor
10 binding to stimulatory ligands or inactivation of the CD28 signal transduction pathway.

The method of immunotherapy of the present invention takes advantage of the surprising and heretofore unappreciated effects of stimulation of the CD28 surface receptor molecule of activated T cells. By activated T cells is
15 meant cells in which the immune response has been initiated or "activated," generally but not necessarily by the interaction of the T cell receptor (TCR)/CD3 T cell surface complex with a foreign antigen or its equivalent. While such activation results in T cell proliferation, it results in only limited induction of T cell effector functions such as lymphokine production.

20 Stimulation of the CD28 cell surface molecule with anti-CD28 antibody results in a marked increase of T cell lymphokine production. Surprisingly, even when the stimulation of the TCR/CD3 complex is maximized, upon costimulation with anti-CD28, there is a substantial increase in the levels of IL-2 lymphokine, although there is no significant increase in T cell proliferation over that induced
25 by anti-TCR/CD3 alone. Even more surprisingly, not only are IL-2 levels significantly increased, but the levels of an entire set of lymphokines, hereinafter referred to as T_H CD28 lymphokines, previously not associated with CD28 stimulation are increased. Remarkably both the T cell proliferation and increased lymphokine production attributable to CD28 stimulation also exhibit resistance to
30 immunosuppression by cyclosporine and glucocorticoids.

The method of immunotherapy of the present invention thus provides a method by which the T cell-mediated immune response can be regulated by stimulating the CD28 T cell surface molecule to aid the body in ridding itself of infection or cancer. The method of the present invention can also be used not

- 4 -

only to increase T cell proliferation, if so desired, but to augment or boost the immune response by increasing the levels and production of an entire set of T cell lymphokines now known to be regulated by CD28 stimulation.

Moreover, because the effectiveness of CD28 stimulation in enhancing
5 the T cell immune response appears to require T cell activation or some form of stimulation of the TCR/CD3 complex, the method of immunotherapy of the present invention can be used to selectively stimulate T cells preactivated by disease or treatment to protect the body against a particular infection or cancer, thereby avoiding the non-specific toxicities of the methods presently used to
10 augment immune function. In addition, the method of immunotherapy of the present invention enhances T cell-mediated immune functions even under immunosuppressed conditions, thus being of particular benefit to individuals suffering from immunodeficiencies such as AIDS.

It will also be appreciated that although the following discussion of the
15 principles of the present invention exemplifies the present invention in terms of human therapy, the methods described herein are similarly useful in veterinary applications.

A better understanding of the present invention and its advantages will be had from a reading of the detailed description of the preferred embodiments
20 taken in conjunction with the drawings and specific examples set forth below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a bar graph illustrating the absence of augmentation of the uptake of thymidine by CD28 stimulated T cells.

Figure 2 is a bar graph illustrating the increase in uridine incorporation by
25 CD28 stimulation of anti-CD3 stimulated T cells.

Figure 3 is a graph illustrating the elevated cyclosporine resistance of T cell proliferation induced by CD28 stimulation.

Figure 4 is a Northern blot analysis of the effects of cyclosporine on PMA-or anti-CD3 activated T cell lymphokine expression induced by anti-CD28.

Figure 5 is a graph illustrating *in vivo* activation of T cells in monkeys by
30 CD28 stimulation.

Figures 6A and 6B are graphs representing changes in lymphocyte levels after infusion of anti-CD28 Mab.

- 5 -

Figures 7A and 7B are graphs representing *in vitro* production of TNF and IL-6 by PBLs under various conditions.

Figures 8A and 8B are graphs representing serum concentration of IL-1 β after single and multiple doses of anti-CD28 Mab.

5 Figures 9A and 9B are graphs representing the serum concentration of IL-6 after single and multiple doses of anti-CD28 Mab.

Figures 10A and 10B are graphs representing IL-6 production of *in vitro* stimulated PBLs isolated from monkeys treated with a single bolus or multiple injections of anti-CD28 Mab.

10 Figure 11 is a graph representing the inhibitory effect of CTLA-4Ig on ³H-thymidine incorporation in a one-way mixed lymphocyte culture.

Figures 12A and 12B are photographs of cardiac allografts to illustrate histopathology.

Figure 13 is a Kaplan-Meier life analysis of cardiac allograft survival after
15 CTLA-4Ig treatment.

Figure 14 is a bar graph illustrating CTLA-4Ig and cyclosporine as synergistic immunosuppressants.

Figure 15 is a bar graph illustrating the effect of herbimycin A on CD28-stimulated IL-2 production.

20 Figure 16 is a bar graph illustrating activation by SEB and anti-CD28 on purified resting T cells in the presence and absence of a blocking Mab to HLA-DR.

Figure 17 is a bar graph illustrating activation by SEB alone or SEB and blocking Mab to HLA-DR in peripheral blood mononuclear cells.

25 Figure 18 is a graph showing *in vitro* long term growth of CD4⁺ peripheral blood T cells propagated with anti-CD3 and anti-CD28.

Figure 19 is a Northern blot analysis of the enhancement of mRNA for IL-2 and TNF- α after costimulation with anti-CD3 and anti-CD28.

Figure 20 is a Northern blot analysis of the ability of mitogens to induce
30 CTLA-4 mRNA expression.

Figure 21 is a Northern blot analysis of the induction of CTLA-4 mRNA expression by costimulation with anti-CD3 Mab and soluble anti-CD28 Mab.

Figure 22 is a graph illustrating the effects on disease progression of CTLA-4Ig treatment of syngeneic, MBP-sensitized cells used to adoptively

- 6 -

transfer the murine autoimmune disease, Experimental Autoimmun Encephalomyelitis (EAE).

Figure 23 is a graph illustrating the effect on disease progression of CTLA-4Ig or control IgG treatment of donor mice and/or isolated cells used to
5 adoptively transfer EAE.

Figure 24 is a graph depicting the effect on disease severity of direct administration of CTLA-4Ig or control human IgG to PLSJLFI/J mice with adoptively transferred EAE.

Figure 25 is a graph illustrating the effect on disease progression of direct
10 administration of CTLA-4Ig or control human IgG to SJL/J mice with adoptively transferred EAE.

Figure 26 is a graph depicting the effect of direct administration of CTLA-4Ig or IgG on disease severity in SJL/J mice with adoptively transferred EAE.

Figure 27 is a graph illustrating the effect on disease severity of direct
15 administration of CTLA-4Ig or control IgG to PLSJLFI/J mice directly immunized with MBP and treated with PT.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In one preferred embodiment of the immunotherapeutic method of the present invention, the CD28 molecule is stimulated to enhance the T cell-
20 mediated immune response of antigen or otherwise activated T cells. CD28 is a 44 kilodalton protein expressed on the surface of about 80% mature T cells which exhibits substantial homology to immunoglobulin genes. See Poggi, A. *et al.*, *Eur. J. Immunol.*, 17:1065-1068 (1987) and Aruffo, A. *et al.*, *PNAS (USA)*, 84:8573-8577 (1987), both herein incorporated by reference. Binding of
25 the CD28 molecule's extracellular domain with anti-CD28 antibodies in accordance with the method of the present invention results in an increase in T cell proliferation and elevated lymphokine levels.

In Specific Examples III-IV and VI-VIII, T cell activation was accomplished by stimulating the T cell TCR/CD3 complex (which mediates the specificity of
30 the T cell immune response) with immobilized anti-CD3 monoclonal antibodies, such as mAb G19-4, or by chemically stimulating with PMA and ionomycin. It should also be appreciated, however, that activation of the T cell can instead be accomplished by routes that do not directly involve CD3 stimulation, such as the stimulation of the CD2 surface protein.

- 7 -

In practice, however, an activated T cell population will be provided by the patient's own immune system, which, barring total immunosuppression, will have T cells activated in response to any foreign or substantially elevated level of antigen present due to disease, infection, inoculation or autoimmunity. The term "foreign antigen" is used broadly herein, meaning an antigen which is either not normally produced by the organism, or, as in carcinomas, an antigen which is not normally produced by the cell which is producing it. The term is also meant to include an antigen which should normally be seen as "self," but, as occurs in autoimmune disease states, provokes an immune response as would a foreign antigen. By "substantially elevated" level of antigen is meant an antigen level exceeding normal ranges and having potentially deleterious effects to the organism due to such elevation.

In accordance with the method of the present invention, stimulation of the CD28 molecule itself is achieved by administration of a ligand, such as a monoclonal antibody or a portion thereof, (e.g. F(ab')₂), having a binding specificity for and stimulatory effect on CD28. Suitable antibodies include mAb 9.3, an IgG2a antibody on deposit with the ATCC which has been widely distributed and is available (for non-commercial purposes) upon request from Dr. Jeffrey A. Ledbetter of Oncogen Corporation, Seattle, WA, or Mab Kolt-2. Both these monoclonal antibodies have been shown to have binding specificity for the extracellular domain of CD28 as described in "Leukocyte Typing II," Ch. 12, pp. 147-156, ed. Reinherz, E. L. *et al.* (1986). Monoclonal antibody 9.3 F(ab')₂ has also been shown to be a satisfactory ligand capable of stimulating the CD28 receptor. It should also be understood that the method of the present invention contemplates the use of chimaeric antibodies as well as non-immunoglobulin, natural and recombinant ligands which bind the CD28 surface molecule. More recently, the natural ligand for CD28, B7/BB1, has also been identified and can be used in accordance with the principles of the present invention. See Linsley, P.S. *et al.*, *J. Exp. Med.* 174:561 (1991). In addition, binding homologs of a natural ligand, whether natural or synthesized by biochemical, immunological, recombinant or other techniques, can also be used in accordance with the principles of the present invention. It will be appreciated that the ligands referred to herein can be utilized in their soluble or cell-bound forms, depending

- 8 -

on their application. Monoclonal antibody 9.3 and B7 are currently preferred stimulatory ligands.

The extracellular domain of CD28, which was sequenced by Aruffo, A. *et al.*, *PNAS (USA)*, 84:8573-8577 (1987), generally comprises the following

5 amino acid sequence:

MetLeuArgLeuLeuLeuAlaLeuAsnLeuPheProSerIleGln
 ValThrGlyAsnLysIleLeuValLysGlnSerProMetLeuVal
 AlaTyrAspAsnAlaValAsnLeuSerCysLysTyrSerTyrAsn
 LeuPheSerArgGluPheArgAlaSerLeuHisLysGlyLeuAsp
 10 SerAlaValGluValCysValValTyrGlyAsnTyrSerGlnGln
 LeuGlnValTyrSerLysThrGlyPheAsnCysAspGlyLysLeu
 GlyAsnGluSerValThrPheTyrLeuGlnAsnLeuTyrValAsn
 GlnThrAspIleTyrPheCysLysIleGluValMetTyrProPro
 ProTyrLeuAspAsnGluLysSerAsnGlyThrIleIleHisVal
 15 LysGlyLysHisLeuCysProSerProLeuPheProGlyProSer
 LysPro

By the term "extracellular domain" as used hereinafter in the specification and claims, is meant the amino acid sequence set forth above, any substantial portion thereof, or any sequence having substantial homology thereto.

20 As shown by the data of Specific Examples III-V, substantial augmentation of the T cell-mediated immunoresponse by CD28 stimulation appears specific for activated T cells. Such specificity is of particular clinical importance and is one of the significant advantages of the method of immunotherapy of the present invention. Administration of anti-CD28
 25 antibodies or other CD28 ligands will specifically augment the response of T cells which are already activated and engaged in the immune response or those in the process of activation. It should, however, also be appreciated that CD28 stimulation may be effective even where the T cells are activated after the binding of the CD28-specific ligand of the present invention to CD28 receptor.
 30 Thus, the T cells at or near the tumor site or site of infection, which are being activated by the antigens produced or present at those sites, will be selectively "boosted" by the CD28 stimulation.

Boosting of the immune response can also be beneficial to healthy individuals, for example, in augmenting their response to antigens presented in

- 9 -

vaccines (see Specific Example IX). CD28 stimulation coupled with antigen administration in accordance with the present invention can result in more effective immunization, not only with conventional vaccines, but in situations where an adequate immune response is difficult to elicit, *e.g.* with human retroviruses such as HIV and some herpes viruses. Examples where CD28 stimulation of the present invention can be used to augment the immune response include, but are not limited to viral vaccines against measles, influenza, polio, herpes viruses (*i.e.* HCMV, Epstein Barr Virus, Herpes Simplex Type I and II); bacterial vaccines against whooping cough (*Bordetella pertussis*), tetanus (*Clostridium tetanus*), pneumonia (*Streptococcus pneumoniae*), meningitis and gonorrhea (*Neisseria*) and against enteropathic bacteria such as *Salmonella*, *E. coli* and *Shigella*. The principles of the present invention are also applicable in inoculations against parasitic infection, including those caused by protozoal parasites, *e.g.* malaria, trypanosomiasis, leishmaniasis, amebiasis, toxoplasmosis, pneumocystis carinni and babesiosis, by cestodes (*e.g.* tapeworm) and by other parasitic organisms. It will also be appreciated that immunization for a humoral response through injection of cDNA for intracellular antigenic production, as described in *Nature* 356:152 (1992), costimulated with anti-CD28 is also contemplated as within the scope of the present invention.

Indeed, recently CD28 engagement and signal transduction have been used to identify IL-13 (Minty, A. *et al.*, *Nature* 362: (6417):248-50 (1993)), a lymphokine involved in the inflammatory and immune response. Punnonen, J. *et al.*, *PNAS (USA)* 90(8): 3730-4 (1993). When TCR/CD3 stimulation is maximized, although T cell proliferation is not markedly increased, the levels of certain lymphokines are substantially increased by CD28 activation, indicating an increase in cellular production of these lymphokines. Thus, in patients undergoing natural maximal TCR/CD3 stimulation or its equivalent T cell activation *in vivo* due to disease or infection, the administration of anti-CD28 antibody or other CD28 ligand to stimulate CD28 in accordance with the method of the present invention will result in substantially elevated lymphokine production.

The increase in lymphokine production achieved by administration of a CD28 stimulator in accordance with the method of the present invention, as particularly shown in Specific Example III, surprisingly results in the increased

- 10 -

production of an entire set of lymphokines, indicating that these lymphokines are under some form of CD28 regulation. Part of this set of lymphokines, which includes IL-2, TNF- α , LT, IFN- γ , and IL-3 as later determined, is somewhat analogous to the T_H1 cell lymphokines present in the mouse which were described by Mosmann, T. R. *et al.*, *Immunol. Today*, 8:223-227 (1987). Although it was originally believed that human IL-4 production was not increased by CD28 stimulation, more recent assays as set forth in Specific Example III have now also shown an increase in the production of other lymphokines, including IL-4 and IL-5 and the increased production of IL-6 and IL-1 was also confirmed in Specific Example IX. It will be appreciated, however, that the term "T_H1 lymphokines" was originally used for ease of reference and was expressly not limited to the lymphokines listed above, but was meant to include all lymphokines whose production is affected or regulated by the binding or stimulation of the CD28 T cell surface molecule. Thus the group of lymphokines affected by CD28 will hereinafter be referred to as T_HCD28 lymphokines, it will again be appreciated that the term T_HCD28 lymphokine is not intended to be limiting to the specific lymphokines listed herein. Furthermore, it will be appreciated that the principles of the present invention can be used in veterinary applications to increase the T cell-mediated immune response and lymphokine production in animals. The term T_HCD28 lymphokines, as used herein, is thus also meant to include analogous animal lymphokines. Thus, by administration of anti-CD28 antibodies or other CD28 ligands in accordance with the method of the present invention, the production and levels of an entire set of lymphokines can be significantly increased.

The method of immunotherapy of the present invention can also be used to facilitate the T cell-mediated immune response in immunodepressed patients, such as those suffering from AIDS. As shown in Specific Examples VI - VIII, T cell proliferation and the increased levels or production of CD28-regulated lymphokines continue to function even in the presence of immunosuppression such as that caused by cyclosporine or dexamethasone. Thus administration of CD28 stimulators such as mAb 9.3 or other CD28 ligands can be used to treat immunodepressed patients to increase their *in vivo* lymphokine levels.

In addition, a variety of syndromes including septic shock and tumor-induced cachexia may involve activation of the CD28 pathway and augmented

- 11 -

production of potentially toxic levels of lymphokines. The immune response can also be deleterious in other situations such as in organ transplant recipients or in autoimmune disease. Thus down-regulation or inactivation of the CD28 pathway, as discussed more fully below and in Specific Examples X and XI, can also provide immunotherapy for those and other clinical conditions.

It should be appreciated that administration of an anti-CD28 antibody has not heretofore been seriously contemplated as a potential immunotherapeutic method for the substantial increase of lymphokine levels at the sites of activated T cells. For example, the addition of mAb 9.3 has been thought only to somewhat augment T cell proliferation, not to induce substantial increases in T_H CD28 lymphokine production.

Although it is not the intent herein to be bound by any particular mechanism by which CD28 binding regulates the T cell-mediated immune response, a model for the mechanism of stimulation has been postulated and supported with experimental data, some of which is shown in Specific Example VIII. It has previously been shown that a number of lymphokine genes undergo more rapid degradation in the cytoplasm than mRNAs from constitutively expressed housekeeping genes, leading to the hypothesis that the instability of these inducible mRNAs has been selected to allow for rapid regulation of gene expression. It is believed that the mechanism of CD28 regulation herein described and claimed is related to the stabilization of rapidly degradable mRNAs for the set of T_H CD28 lymphokines set forth above. To date, it appears no other mechanism in any eukaryotic cell system has been described to demonstrate that a cell surface activation pathway can alter gene expression by inducing specific alteration in mRNA degradation. (A more in-depth analysis of possible models of CD28 activation is presented later herein.)

As shown in Specific Example IV, costimulation of CD28 and TCR/CD3 caused an increase in mRNA of the T_H CD28 lymphokines which was not the result of a generalized increase in a steady state mRNA expression of all T cell activation-associated genes. The increase was disproportionate and thus could not be accounted for by the increase in percentage of proliferating cells in culture. These data, in addition to further studies not detailed herein, demonstrate that activation of the CD28 surface molecule of activated T cells functions to specifically stabilize lymphokine mRNAs. Increased mRNA stability,

i.e. slower degradation thereof, results in increased translation of the mRNA, in turn resulting in increased lymphokine production per cell. An increase in per T cell production of lymphokines that allows the T cell to influence the response of other inflammatory and hematopoietic cells is referred to as paracrine production. In contrast, an increase in lymphokine levels merely due to increased cell proliferation, such as that shown in Martin, P.J. *et al.*, *J. Immunol.* 136:3282-3287 (1986), is commonly referred to as autocrine production. For ease of reference, paracrine production is also herein referred to as "cellular" production of lymphokines.

Thus, in accordance with the principles of the present invention, ligands with binding specificity for the CD28 molecule are administered in a biologically compatible form suitable for administration *in vivo* to stimulate the CD28 pathway. By "stimulation of the CD28 pathway" is meant the stimulation of the CD28 molecule resulting in increased T cell production of T_HCD28 lymphokines. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the ligand to be administered in which the toxic effects, if any, are outweighed by the therapeutic effects of the ligand. Administration of the CD28 ligand can be in any suitable pharmacological form, including but not limited to intravenous injection of the ligand in solution.

It should be understood that, although the models for CD28 regulation of lymphokine production are described with respect to stimulation and enhancement of lymphokine levels, as noted above, down-regulation or inhibition of the CD28 pathway is also in accordance with the principles of the present invention. Down-regulation or suppression of the immune response is of particular clinical interest for a variety of conditions, including septic shock, tumor-induced cachexia, autoimmune diseases and for patients receiving heart, lung, kidney, pancreas, liver and other organ transplants. One preferred approach to down-regulation is the blocking of the CD28 receptor stimulatory binding site on its natural ligand. For example, CTLA-4 (discussed in more detail below), which shares 32% amino acid homology with CD28 and appears to have greater binding affinity for B7 than CD28, can be used to bind B7 and prevent CD28 binding and activation thereby. See Linsley, P.S. *et al.*, *J. Exp. Med.*, 174:561 (1991). Such regulation has been accomplished *in vivo* as described in Specific Example X. In this Example, acute rejection of fully MHC-

- 13 -

mismatched cardiac allografts was prevented by blocking B7-dependent T cell activation, *i.e.* CD28 binding, with the soluble recombinant fusion protein CTLA-4Ig. The immunosuppression observed with CTLA-4Ig did not result in significant alterations in circulating T cell subsets. It will be appreciated that
5 other B7-binding ligands such as a monoclonal antibody to B7 can be similarly employed. In addition, CTLA-4Ig treatment can be efficacious in the treatment of autoimmune diseases, as shown in the murine model for multiple sclerosis (*i.e.* Experimental Autoimmune Encephalomyelitis (EAE)). CTLA-4Ig treatment of T cells isolated from a mouse immunized with Myelin Basic Protein (MBP) results
10 in reduced disease severity when the treated cells are introduced into a syngeneic animal. Likewise, when mice immunized with MBP and injected with pertussis toxin (PT) are treated directly with CTLA-4Ig, disease severity is reduced. These findings confirm the *in vivo* immunosuppressive effects of CTLA-4Ig treatment. Thus, administration of CTLA-4 can provide an effective
15 therapeutic strategy for the treatment of autoimmune diseases. Those skilled in the art will also appreciate that the cell lines and animal models used to exemplify the present invention are recognized predictors of efficacy in humans.

It will be appreciated that down-regulation can also be accomplished by blocking CD28 receptor binding to B7 by occupying the CD28 binding site with
20 nonstimulatory ligands which may mimic stimulatory ligands but do not result in activation of the CD28 pathway, *e.g.* Fabs, modified natural, synthetic, recombinant or other ligands which do not crosslink or otherwise do not activate receptors.

As discussed above and shown in the Specific Examples, the blockade of
25 stimulatory ligands which bind to CD28 and activate the CD28 pathway (*e.g.* B7) or the blocking of the CD28 binding site can reduce the increased lymphokine expression which occurs upon CD28 activation. Thus, just as manipulation of the CD28 pathway can be used to enhance T cell immune responses, it can also be used to suppress such responses. Since unregulated
30 lymphokine production has been implicated in the aetiology of autoimmunity, CD28-mediated immunosuppression can be exploited to treat various autoimmune diseases. Methods of suppressing the CD28 pathway in accordance with the present invention are desirable since this pathway is resistant to the effects of cyclosporine, which is commonly used as an

- 14 -

immun suppressive agent in the treatment of autoimmune diseases.

Immunosuppression via the CD28 pathway can restore immunoregulation and thus reduce the pathologic effects of such autoimmune diseases as systemic lupus erythematosus, rheumatoid arthritis, hemolytic anemia, myasthenia gravis, 5 scleroderma, Sjögren's syndrome, ulcerative colitis, multiple sclerosis, and a host of other systemic as well as organ-specific autoimmune diseases.

Administration of stimulatory ligands (e.g. mAb 9.3, Kolt, B7) or inhibitory ligands (e.g. CTLA-4lg, Fab fragments of mAb 9.3, and the like) can be in any suitable pharmacological form, including parenterally or topically.

10 Pharmaceutical compositions may also take the form of ointments, gels, pastes, creams, sprays, lotions, suspensions, solutions and emulsions of the active ingredient in aqueous or nonaqueous diluents, syrups, granulates or powders. In addition to a ligand of the present invention, the pharmaceutical compositions can also contain other pharmaceutically active compounds or a plurality of 15 compounds of the invention.

More particularly, a ligand of the present invention (also referred to herein as the active ingredient) may be administered by any suitable route, including parenteral (including subcutaneous, intramuscular, intravenous and intradermal), topical (including transdermal, buccal and sublingual), rectal, 20 vaginal, nasal and pulmonary. Although the preferred route of administration is currently parenteral, it will be appreciated that the delivery route of choice will vary with the condition and age of the recipient and the nature of the disease or condition being treated.

While it is possible for the active ingredient to be administered alone, it is 25 preferable to present it as a pharmaceutical formulation comprising at least one active ingredient, as defined above, together with one or more pharmaceutically acceptable carriers therefor and optionally other therapeutic agents. Each carrier must be "acceptable" in the sense of being compatible with other ingredients of the formulation and not injurious to the patient.

30 Formulations suitable for parenteral administration include aqueous and non-aqueous isotonic sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents, and

- 15 -

liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring
5 only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

Formulations include those suitable for parenteral (including
10 subcutaneous, intramuscular, intravenous and intradermal), topical (including transdermal, buccal and sublingual), rectal, vaginal, nasal and pulmonary administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods include the step of bringing into association the active ingredient
15 with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into associated the active ingredient with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

Pharmaceutical compositions for topical administration according to the
20 present invention may be formulated as an ointment, cream, suspension, lotion, solution, paste, gel, spray aerosol or oil. Alternatively, a formulation may comprise a patch or a dressing such as a bandage or adhesive plaster impregnated with active ingredients and optionally one or more excipients or diluents. Formulations suitable for topical administration in the mouth include
25 mouthwashes comprising the active ingredient in a suitable liquid carrier. It will also be appreciated that in a carrier suitable to preserve efficacy of the ligand, oral administration is also contemplated.

For conditions of the eye or other external tissues, *e.g.* mouth and skin, the formulations are preferably applied as a topical ointment or cream containing
30 the active ingredient. When formulated in an ointment, the active ingredient may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the

- 16 -

skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogues. Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient.

Formulations for rectal administration may be presented as a suppository with a suitable base. Formulations for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient, such carriers as are known in the art to be appropriate. Formulations suitable for nasal administration, include formulations wherein the carrier is a liquid for administrations, for example, a nasal spray or a nasal drops, include aqueous or oily solutions of the active ingredient.

Preferred unit dosage formulations are those containing a daily dose or unit, daily subdose, or an appropriate fraction thereof, of the active ingredient. In any event, in practicing the present invention, the amount of active ingredient to be used or administered, alone or in combination with other agents, will vary with the patient being treated and will be monitored on a patient-by-patient basis by the physician. Generally, a therapeutically effective amount of the vaccine will be administered for a therapeutically effective duration. By "therapeutically effective amount" and "therapeutically effective duration" are meant an amount and duration to achieve the result desired in accordance with the present invention without undue adverse or with medically acceptable physiological effects, which effects can be determined by those skilled in the medical arts. It will also be appreciated that, particularly when a natural ligand is used in the practice of the invention, its isolation or production should render it substantially free of undesirable contaminants such as other proteins (*i.e.* "substantially pure") which may adversely impact on its efficacy or use. Acceptable levels of purity can be determined by those skilled in the pharmaceutical and medical arts and can depend on the specific ligand and composition and its intended use.

It should also be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. In accordance with the present invention, ligands may also be

- 17 -

presented for the use in the form of veterinary formulations, which may be prepared, by methods conventional in the art.

Accumulating evidence suggests that in addition to T cell receptor occupancy, other costimulatory signals are required to induce a complete T cell-mediated immune response. The CD28 receptor expressed on T cells serves as a surface component of a novel signal transduction pathway that can induce paracrine levels of cellular production of lymphokines. Interaction of CD28 with its natural ligand B7 which is expressed on the surface of activated B cells macrophages or dendritic cells can act as a costimulus to induce high level lymphokine production in antigen receptor-activated T cells. Thus, another approach to down-regulation is to inhibit the activation of the CD28 signal transduction pathway as described below.

Although the CD28 signal transduction pathway is not entirely understood, Specific Example XI demonstrates that binding of CD28 induces protein tyrosine phosphorylation distinct from T cell receptor (TCR)-induced tyrosine phosphorylation. For example, TCR-induced tyrosine phosphorylation occurs in both resting and activated T cells, while CD28-induced tyrosine phosphorylation occurs primarily in previously activated T cells. Most striking were the results after CD28 receptor ligation by cell-bound B7, where phosphorylation was consistently detectable on only a single substrate. Experiments using the Jurkat E6-1 T cell line indicated an absolute requirement for PMA pretreatment in order to observe CD28-induced tyrosine phosphorylation. In contrast, there was no requirement for cellular preactivation in the Jurkat J32 line, while, as noted above, there was a relative requirement for PMA or TCR prestimulation of normal T cells in order to induce CD28 responsiveness. Studies with Jurkat mutants further indicated that CD28-induced tyrosine phosphorylation and biologic function can occur in the absence of the TCR. In this respect, CD28 appears to be unique in that other accessory molecules involved in T cell activation, such as CD2, Ly-6, Thy-1, and CD5 appear to require the presence of the TCR. Thus, specific tyrosine phosphorylation appears to occur directly as a result of CD28 ligand binding and is involved in transducing the signal delivered through CD28 by accessory cells that express the B7/BB1 receptor.

- 18 -

Studies with an inhibitor of the *src* family of tyrosine kinases, herbimycin A, and with tyrosine phosphatase, as described in Specific Example XI, further show that the functional effects of CD28 stimulation on lymphokine gene expression require the above-described protein tyrosine phosphorylation. The data on tyrosine phosphorylation inhibitors thus demonstrate that inactivation of CD28-mediated signal transduction can also be used to down-regulate lymphokine production in accordance with the principles of the present invention.

Immunotherapy through CD28 stimulation in accordance with the present invention also has clinical applicability in the treatment of bone marrow transplant recipients. The success of autologous and allogeneic bone marrow transplantation (BMT) has generally been limited by recurrent malignancy, graft-vs-host disease (GVHD), and the life-threatening immune deficiency that occurs after BMT. One approach to overcoming these problems has been the adoptive transfer of lymphocytes in combination with lymphokine infusions, to accelerate immune reconstitution or mediate cytotoxicity directed at malignant cells. The side effects attending such transfer with lymphokine infusions are, however, quite significant.

Thus, T cell proliferation and lymphokine synthesis in the absence of exogenously added IL-2 in response to CD3 and CD28 costimulation, as described in Specific Example XIII, provides a unique opportunity for clinical use of adoptive transfer of activated T cells to repair T cell defect *in vivo* without exogenous lymphokine infusions. The studies detailed in Specific Example XIV show that defective *in vitro* proliferative responses to anti-CD3 (OKT3 or G19-4) can be repaired by adding mAb 9.3 to the cultures. See Joshi, I. *et al.*, *Blood Suppl.* (Abstract) (1991). Costimulation of T cells with OKT3/9.3 repaired proliferative responses as a result of increasing the levels of mRNA expression for cytokines/lymphokines such as IL-2, GM-CSF, and TNF α . See Joshi, *supra*; Perrin, P.J. *et al.*, *Blood Suppl.* (Abstract) (1991). Purified normal CD4⁺ cells can thus be costimulated with OKT3/9.3 to expand and secrete lymphokines for long periods of time. Preclinical studies using mAb 9.3 stimulation have shown no untoward effects in monkeys. Although CD3-stimulated cells (CTC) provide helper factors to normal B cells, CD3-CD28 costimulated cells appear even more effective in producing helper factors than CD3-stimulated T cells. Thus,

- 19 -

costimulation may also enhance the growth of helper cells and cytotoxic T cells for adoptive immunotherapy after BMT. The administration *in vivo* of T cells that have been expanded *in vitro*, will provide two prominent benefits in marrow transplantation. The ability of CD28- treated cells to produce many lymphokines
5 which have a strong positive effect on hematopoiesis, such as GM-CSF, IL-3, and IL-6, should accelerate engraftment after marrow transplantation. In addition, the ability of anti-CD28 to trigger cytotoxicity and to cause the production of lymphokines such as TNF is a novel form of adoptive immunotherapy that should augment the anti-neoplastic efficacy of bone marrow
10 transplantation.

The possible role of CD28 in energy was also examined. Generally, the activation of a quiescent T cell is initiated through stimulation of the T cell antigen receptor. This activation can occur either through engagement of an antigenic peptide presented in the antigen binding groove of a self-encoded MHC
15 molecule or by engagement of a foreign MHC molecule. However, while this receptor-mediated activation event is required for the initiation of a T cell response in a quiescent cell, recent studies have demonstrated that signals transduced by the antigen receptor alone are not sufficient to lead to an effective T cell-mediated immune response. Several *in vitro* models suggest
20 that, in fact, T cell receptor (TCR)/CD3 activation alone of a quiescent T cell leads to the induction of a state in which the T cell becomes anergic to further stimulations through its antigen-specific receptor. This state is relatively long-lived and, for at least several weeks, renders that cell incapable of further response upon antigenic stimulation. It is hypothesized that this isolated
25 activation of the TCR/CD3 complex alone in the absence of additional T cell costimulatory molecules plays an important role in regulating a peripheral immune response by preventing T cells from responding to self antigens in the periphery.

Thus, for T cells to mount a proliferative response and initiate a cell-
30 mediated immune response, a quiescent T cell normally requires stimulation not only through its antigen-specific T cell receptor but also through a second receptor which provides additional costimulatory signals to the cell. The data set forth herein, *e.g.* in Specific Example XII, demonstrates that CD28 provides an essential costimulatory signal for T cell responses *in vitro* and *in vivo*. Thus,

- 20 -

the CD28 receptor's ability to augment T cell lymphokine production not only results in the initiation of a cell-mediated immune response, but also prevents the induction of anergy in a quiescent T cell.

The role of CD28 in the prevention of programmed cell death has also
5 been tested. The induction of cell death has a major role in the elimination of self-reactive or non-reactive T cells in the thymus. In the thymus, it is thought that T cell receptor signals are able to induce programmed cell death, in a selective and specific fashion involving cells that express T cell receptors specific for self-antigens. As described in Turka, L.A. *et al.*, *J. Immunol.*,
10 144:1646-53 (1990), CD28 is expressed in developing T cells in the thymus, and the binding of mAb 9.3 prevents thymocyte cell death. Programmed cell death is also thought to occur in mature T cells in peripheral lymphoid organs. Signals delivered through the T cell receptor can induce cell death (see Newell, M.K., *et al.*, *Nature*, 347:286-8 (1990)). It has also been proposed that cell
15 death may have a role in certain forms of immunopathology. For example, in HIV-1 infection it has been proposed that the progressive immunodeficiency may be the result of immunologically-mediated cell death, rather than a direct consequence of viral-induced cytopathic effects. See Ameisen, J.C. *et al.*, *Immunol. Today*, 4:102 (1991); also see Groux H., *et al.*, *J. Exp. Med.*
20 175:331-340 (1992). The results in Specific Example XIII show that CD28 can prevent cell death in mature T cells. Thus, abnormal expression or activation of CD28 may have a role in immunopathology of certain autoimmune disorders such as systemic lupus erythematosus, a disorder characterized by abnormally self-reactive T cells that have failed to undergo elimination in the thymus or
25 escape from anergic states in the peripheral lymphoid system. Similarly, the ability to induce CD28 activation may be beneficial in disorders characterized by progressive T cell depletion such as HIV-1 infection.

It was also demonstrated in Specific Example XII that superantigens SEA and SEB, which do not require traditional processing for binding to MHC, can
30 directly activate purified T cells in the absence of accessory cells as determined by a transition from G₀ to G₁ and induction of IL-2 receptor expression. However, neither SEA nor SEB alone was sufficient to result in T cell proliferation. The induction of T cell proliferation by SEB or SEA required the addition of a second costimulatory signal. This could be provided by either

- 21 -

accessory cells or monoclonal antibody stimulation of CD28. As previously reported, T cell proliferation induced by enterotoxin in the presence of accessory cells was partially inhibited by a blocking antibody (HLA-DR) against class II MHC. In contrast, in purified T cells when costimulation was provided through CD28, proliferation was not inhibited by class II antibody and HLA-DR expression was not detectable. In addition, costimulation through CD28 was partially resistant to the effects of cyclosporine. These results demonstrate that CD28 costimulation is sufficient to induce lymphokine production and subsequent proliferation of enterotoxin-activated T cells, and that this effect is independent of class II MHC expression. This prevention of *in vivo* CD28 activation of superantigen-activated cells such as those occurring in toxic shock syndrome and rheumatoid or Lyme arthritis, may substantially decrease disease morbidity.

Although as noted previously the present invention is not intended to be restricted to specific mechanisms of activation, the role of CTLA-4 in the CD28 activation pathway has been examined and models consistent with the data presented herein have been postulated. (See *e.g.* Specific Example XV.) Recent work in our laboratory has shown that the CTLA-4 gene lies immediately adjacent to CD28 on chromosome 2, with a similar genomic organization and 32% amino acid homology. Based on their chromosomal localization and sequence and organizational similarities, CD28 and CTLA-4 likely represent evolutionary gene duplication. By standard nomenclature they might thus more appropriately be named CD28 α and CD28 β , although the terms CD28 and CTLA-4 are retained herein.

Although CTLA-4 is not expressed on quiescent lymphoid cells, its expression at the RNA level can be rapidly induced upon T cell activation. Two potential mechanisms by which CTLA-4 might function are postulated as follows. First, since CD28 and CTLA-4 contain an unpaired cysteine in their extracellular domain, this cysteine residue may be used to form crosslinked dimeric receptors on the surface. If this were the case, it may suggest that CTLA-4 is normally expressed on the surface as a heterodimer with CD28. Under such conditions, the higher affinity of CTLA-4 for the natural ligand B7 might in the dimeric state lead to a higher affinity receptor with enhanced signaling capabilities. This might allow for an enhanced signal transduction

- 22 -

capability through the CD28-CTLA-4 heterodimer in an antigen-activated cell. In addition, if CD28 and CTLA-4 are found primarily in activated cells in a heterodimeric state, this might account for observations that CD28-containing receptors have enhanced signaling capabilities in activated cells.

5 On the other hand, the data presented herein are also compatible with a model in which CTLA-4 is induced upon T cell activation as a competitive inhibitor of CD28 and is used to down-modulate an ongoing immune response by inhibiting further interactions between B7 and CD28 on the surface. In addition, it is quite possible that the CTLA-4 expressed on the surface is also
10 expressed in a shed form, and this shed form of the receptor acts as a soluble competitive inhibitor of an ongoing B7-CD28 interaction, thereby preventing the antigen-presenting cell from activating additional T cells in its environment. Thus, the ability of T cells to produce an additional isoform of CD28, *i.e.* CTLA-4, suggests that the interplay of expression of CD28 and CTLA-4 has profound
15 effects on the ability of T cells to be activated through a CD28-containing receptor.

 CD28 pathway activation and inhibition studies indicate that the ability of the CD28 natural ligand B7 to activate a T cell to augment lymphokine production is entirely mediated through a CD28-containing receptor, either a
20 CD28 homodimer or a CD28-CTLA-4 heterodimer. Thus, the data suggest that a CTLA-4 homodimer is not critical in T cell activation, but may play an important role in down-modulation of T cell lymphokine production, while a CD28-CTLA-4 heterodimer may account for the enhanced signaling properties of CD28-containing receptors upon T cell activation.

25 The role of CTLA-4 in CD28-mediated signal transduction event may explain why the novel and profound effects of CD28 on normal T cell activation encountered and described herein were not previously observed in human T cell lines. Earlier work on the CD28 pathway occurred in cell lines such as Jurkat human T cells. Extensive attempts to stimulate these cells to express CTLA-4
30 have been entirely negative (see Figure 20). In contrast, standard activation events that lead to cell cycle progression of normal T cells either through chemical mitogens such as phytohemagglutinin (PHA) and phorbol myristate acetate (PMA) leads to rapid induction of CTLA-4 expression as does crosslinking of the TCR/CD3 complex. Therefore, the inability of previous

- 23 -

investigators to appreciate or harness the CD28 activation pathway to enhance cellular production of lymphokines was likely due to the lack of expression of the CTLA-4 isoform of CD28 in these cell lines. Interestingly, costimulation of resting T cells with anti-CD28 monoclonal antibodies enhances the expression of the CTLA-4 gene (see Figure 21). Thus, the CD28 activation pathway in normal cells may in fact involve a positive feedback loop in which initial CD28 stimulation through the CD28 homodimer enhances the expression of CTLA-4 thus leading to enhanced heterodimer expression and signal transduction. Alternatively, the enhanced CTLA-4 may lead to the production of a receptor which competes for CD28 signal transduction thus leading to the ultimate termination of lymphokine production and acts as a negative feedback loop to down modulate an ongoing CD28-mediated lymphokine production.

SPECIFIC EXAMPLE I

Preparation of CD28 Stimulator Monoclonal Antibody 9.3.

The monoclonal antibody (mAb) 9.3, an IgG2a monoclonal antibody which binds to the extracellular domain of the CD28 molecule, was produced by a hybrid cell line originally derived as described by Hansen *et al.*, *Immunogen.*, 10:247-260 (1980). Ascites fluid containing high titer monoclonal antibody 9.3 was prepared by intraperitoneal inoculation of $5 - 10 \times 10^6$ hybrid cells into a Balb/C x C57BL/6 F₁ mice which had been primed intraperitoneally with 0.5 ml of Pristane (Aldrich Chemical Co., Milwaukee, WI). The monoclonal antibody 9.3 was purified from ascites fluid on a staphylococcal protein-A sepharose column as described by Hardy, R., "Handbook of Experimental Immunology," Ch. 13 (1986).

Prior to use in functional assays, purified mAb 9.3 was dialyzed extensively against phosphate buffered saline (KCl 0.2 grams/liter dH₂O; KH₂PO₄ 0.2 grams/liter dH₂O; NaCl 8.0 grams/liter dH₂O; Na₂HPO₄·7H₂O 2.16 grams/liter dH₂O) and then filtered through a 0.22 cubic micron sterile filter (Acrodisc, Gelman Sciences, Ann Arbor, MI). The mAb 9.3 preparation was cleared of aggregates by centrifugation at 100,000 xg for 45 m at 20°C. The resulting purified mAb 9.3 was resuspended in phosphate buffered saline to a final concentration of 200 µg/ml as determined by OD₂₈₀ analysis and stored at 4°C prior to use.

- 24 -

SPECIFIC EXAMPLE II

Isolation of CD28⁺ T Cells.

Buffy coats were obtained by leukapheresis of healthy donors 21 to 31 years of age. Peripheral blood lymphocytes (PBL), approximately 2.5×10^9 ,
5 were isolated from the buffy coat by Lymphocyte Separation Medium (Litton Bionetics, Kensington, MD) density gradient centrifugation. The CD28⁺ subset of T cells was then isolated from the PBL by negative selection using immunoabsorption, taking advantage of the reciprocal and non-overlapping distribution of the CD11 and CD28 surface antigens as described by Yamada
10 *et al.*, *Eur. J. Immunol.*, 15:1164-1688 (1985). PBL were suspended at approximately 20×10^6 /ml in RPMI 1640 medium (GIBCO Laboratories, Grand Island, NY) containing 20mM HEPES buffer (pH 7.4) (GIBCO Laboratories, Grand Island, NY), 5mM EDTA (SIGMA Chemical Co., St. Louis, MO) and 5% heat-activated human AB serum (Pel-Freez, Brown Deer, WI). The cells were
15 incubated at 4°C on a rotator with saturating amounts of monoclonal antibodies 60.1 (anti-CD11a) (see Bernstein, I.D. *et al.*, "Leukocyte Typing II," Vol. 3, pp. 1-25, ed. Reinherz, E. L. *et al.*, (1986); 1F5 (anti-CD20) (see Clark, E. A. *et al.*, *PNAS(USA)*, 82:1766-1770 (1985)); FC-2 (anti-CD16) (see June, C. H. *et al.*, *J. Clin. Invest.*, 77: 1224-1232 (1986)); and anti-CD14 for 20 m. This mixture of
20 antibodies coated all B cells, monocytes, large granular lymphocytes and CD28⁺ T cells with mouse immunoglobulin. The cells were washed three times with PBS to remove unbound antibody, and then incubated for 1 h at 4°C with goat anti-mouse immunoglobulin-coated magnetic particles (DynaI, Inc., Fort Lee, NJ) at a ratio of 3 magnetic particles per cell. Antibody-coated cells that were
25 bound to magnetic particles were then removed by magnetic separation as described by Lea, T. *et al.*, *Scan. J. Immunol.*, 22:207-216 (1985). Typically, approximately 700×10^6 CD28⁺ T cells were recovered. Cell purification was routinely monitored by flow cytometry and histochemistry. Flow cytometry was performed as described by Ledbetter, J. A. *et al.*, "Lymphocyte Surface
30 Antigens," pp. 119-129 (ed. Heise, E., 1984). Briefly, CD28⁺ T cells were stained with fluorescein isothiocyanate (FITC)-conjugated anti-CD2 mAb OKT11 (Coulter, Hialeah, FL) and with FITC-conjugated anti-CD28 mAb 9.3 as described by Goding, J. W., "Monoclonal Antibodies Principles and Practice," p. 230 (ed. Goding, J. W., 1983). CD28⁺ T cells were over 99% positive with FITC-

- 25 -

- conjugated monoclonal antibody OKT11 and over 98% positive FITC-conjugated monoclonal antibody 9.3 when compared to a non-binding, isotype-matched, FITC-labeled control antibody (Coulter, Hialeah, FL). Residual monocytes were quantitated by staining for non-specific esterase using a commercially available
- 5 kit obtained from Sigma Chemical Co., St. Louis, MO, and were less than 0.1% in all cell populations used in this study. Viability was approximately 98% as measured by trypan blue exclusion as described by Mishell, B.B. *et al.*, "Selected Methods in Cell. Immunology," pp. 16-17 (1980).

SPECIFIC EXAMPLE III

10 Increased Cellular Production of Human T_H CD28 Lymphokines by CD28 Stimulation by Monoclonal Antibody 9.3.

A. Increased Production of IL-2, TNF- α , IFN- γ and GM-CSF.

- CD28⁺ T cells were cultured at approximately 1×10^5 cells/well in the presence of various combinations of stimulators. The stimulators included
- 15 phorbol myristate acetate (PMA) (LC Services Corporation, Woburn, MA) at 3 ng/ml conc.; anti-CD28 mAb 9.3 at 100 ng/ml; anti-CD3 mAb G19-4 at 200 ng/ml which was immobilized by adsorbing to the surface of plastic tissue culture plates as previously described by Geppert, *et al.*, *J. Immunol.*, 138:1660-1666 (1987); also Ledbetter, *et al.*, *J. Immunol.*, 135: 2331-2336
- 20 (1985); ionomycin (Iono) (Calbiochem., San Diego, CA) at 100 ng/ml. Culture supernatants were harvested at 24 h and serial dilutions assayed for the presence of T_H CD28 lymphokines.

- Specifically, IL-2 was assayed using a bioassay as previously described by Gillis *et al.*, *Nature*, 268:154-156 (1977). One unit (U) was defined as the
- 25 amount of IL-2 needed to induce half maximal proliferation of 7×10^3 CTLL-2 (a human cytotoxic T cell line) cells at 24 h of culture. In separate experiments, the relative levels of IL-2 for each of the culture conditions above were independently confirmed using a commercially available ELISA assay (Genzyme Corp., Boston, MA). TNF- α /LT levels were measured using a semi-automated
- 30 L929 fibroblast lytic assay as previously described by Kunkel *et al.*, *J. Biol. Chem.*, 263:5380-5384 (1988). Units of TNF- α /LT were defined using an internal standard for TNF- α (Genzyme Corp., Boston MA). The independent presence of both TNF- α and LT was confirmed by the ability of a monoclonal antibody specific for each cytokine to partially inhibit cell lysis mediated by the

- 26 -

supernatant from cells costimulated with immobilized anti-CD3 mAb G19-4 and anti-CD28 mAb 9.3. IFN- γ was measured by radioimmunoassay using a commercially available kit (Centocor, Malvern, PA). Units for IFN- γ were determined from a standard curve using ^{125}I -labeled human IFN- γ provided in the test kit. GM-CSF was detected by stimulation of proliferation of the human GM-CSF-dependent cell line AML-193, as described by Lange *et al.*, *Blood*, 70:192-199 (1987), in the presence of neutralizing monoclonal antibodies to TNF- α and LT. The ^3H -thymidine uptake induced by 10 ng/ml of purified GM-CSF (Genetics Institute, Cambridge, MA) was defined as 100 U. Separate aliquots of cells were recovered 48 h after stimulation and assayed for the percentage of cells in late stages of the cell cycle (S + G₂ + M) by staining of cells with propidium iodide and analysis by flow cytometry as previously described by Thompson *et al.*, *Nature*, 314:363-366 (1985).

As shown in Table 1, CD28 stimulation of CD3-stimulated T cells resulted in marked increases in cellular production of IL-2, TNF- α , IFN- γ and GM-CSF.

- 27 -

TABLE 1

Increased Cellular Production of T_H CD28 Lymphokines by CD28 Stimulation

	STIMULUS	IL-2	TNF- α /LT	IFN- γ	GM-CSF	S + G ₂ + M
		(U/ml)	(U/ml)	(U/ml)	(U/ml)	(%)
5	Medium	<2	0	0	0	4.6
	PMA	<2	0	0	NT	5.5
	anti-CD28	<2	5	0	0	6.5
	anti-CD28 + PMA	435	300	24	150	48.9
	anti-CD3 ⁱ	36	50	24	120	39.7
10	anti-CD3 ⁱ + anti-CD28	1200	400	74	1050	44.7
	Ionomycin	<2	0	0	NT	6.6
	Ionomycin + PMA	200	5	37	NT	43.6
	Ionomycin + PMA + anti-CD28	1640	320	128	NT	43.5

15

ⁱ = immobilized

NT = not tested

B. Effects of Anti-CD28 Stimulation on T cell IL-4, IL-5 and IL-3

20 Secretion.

Previous studies of the effects of anti-CD28 Mab stimulation on T cell production of lymphokines of the T_H 2 type were limited to the first few days of stimulation. In those studies IL-4 could not be detected after anti-CD28 stimulation (noted in Thompson *et al.*, *PNAS* 86:1333 (1989)). On

25 reexamination of this question, it was found that anti-CD28 can augment production of IL-4 and of IL-5, and thus, augments production of T_H 2 lymphokines as well as the T_H 1 type lymphokines previously described. As can be seen in Table 2, small amounts of both IL-4 and IL-5 can be detected after 24 h of stimulation with anti-CD3 plus anti-CD28. However, when cells are

30 restimulated after 8 days in culture, large amounts of both IL-4 and IL-5 are secreted. As shown in Table 3, similar results were found when the CD28 subset of T cells were stimulated with the combination of phorbol myristate acetate (PMA) and anti-CD28 mAb. These results indicated that small amounts of IL-4 and IL-5 can be detected after initial stimulation of resting T cells with

- 28 -

anti-CD28. However, with continued stimulation, differentiation occurs, and large amounts of IL-4 and, particularly, of IL-5 are produced, while lesser amounts of IL-2 and γ -IFN are also produced.

TABLE 2

5 Effects of anti-CD3 and anti-CD28 Treatment of IL-4 and IL-5 Production by T cells

STIMULUS	INITIAL				RESTIMULATION			
	IL-4 pg/ml	IL-5 pg/ml	IFN pg/ml	IL-2 U/ml	IL-4 pg/ml	IL-5 pg/ml	IFN pg/ml	IL-2 U/ml
Medium	<1	<1	<1	<0.1	<1	<1	<1	<0.1
anti-CD3	<1	<1	630	2.3	220	225	246	0.02
anti-CD3 + anti- CD28	207	137	887	7.0	498	2545	379	0.2

The data in Table 2 above were obtained using the following protocol: CD28⁺ T cells were isolated by negative selection using monoclonal antibodies and magnetic immunobeads as described in Specific Example II. The cells were cultured at 1×10^6 /ml in RPMI medium containing 10% FCS (Medium), or in culture wells containing anti-CD3 monoclonal antibody G19-4 adsorbed to the plastic, or plastic adsorbed anti-CD3 plus anti-CD28 mAb 9.3 added in solution at 0.5 μ g/ml. Supernatants from the cell culture were analyzed for lymphokine concentration using commercially available ELISA kits and the values expressed as pg/ml for IL-4, IL-5 and γ -IFN, or as units per ml, for IL-2. Supernatants were analyzed after 24 h of culture (Initial) or alternatively, the cells were cultured for 8 days in the above forms of stimulation, the cells were recovered, washed, and then restimulated with their original treatment, and the supernatants analyzed after a further 24 h of stimulation (Restimulation). The values represent means of duplicate cultures.

TABLE 3

Effects of PMA and anti-CD28 Treatment on IL-4 and IL-5 Production by T cells

5	STIMULUS	INITIAL				RESTIMULATION			
		IL-4 pg/ml	IL-5 pg/ml	IFN pg/ml	IL-2 U/ml	IL-4 pg/ml	IL-5 pg/ml	IFN pg/ml	IL-2 U/ml
	Medium	<1	<1	<1	<0.1	<1	<1	<1	<0.1
	PMA	<1	<1	152	.07	<1	<1	<1	0.6
10	anti-CD28 + PMA	<1	187	558	8.1	285	392	335	10.3

The data in Table 3 above were obtained using the following protocol: CD28⁺ T cells were isolated by negative selection using monoclonal antibodies and magnetic immunobeads as described in Specific Example II. The cells were cultured at 1×10^6 /ml in RPMI medium containing 10% FCS (Medium), or in medium plus PMA 3 ng/ml, or in PMA plus anti-CD28 Mab 9.3 at 0.5 μ g/ml. Supernatants from the cell culture were analyzed for lymphokine concentration using commercially available ELISA kits and the values expressed as pg/ml for IL-4, IL-5 and γ -IFN. Supernatants were analyzed after 24 h of culture (Initial) or, alternatively, the cells were cultured for 8 days in the above forms of stimulation, and then restimulated, and the supernatants analyzed after a further 24 h period of stimulation (Restimulation). The values represent means of duplicate cultures.

Induction of IL-3 expression in T cells after anti-CD28 treatment. IL-3 is a multilineage hematopoietic growth factor that is primarily produced by T cells, and is generally considered to be produced by T_H1 cells. The experimental protocol and the findings described herein are described in detail in Guba, S.C. *et al.*, *J Clin. Invest.* 84(6):1701-1706 (1989), incorporated herein by reference.

PBL were isolated as described previously. The CD28⁺ subset of T cells was isolated by negative selection as described by June, C.H. *et al.*, *Mol. Cell. Biol.* 7:4472-4481 (1987). In some experiments, the CD28⁺ subset of T cells was isolated by incubating PBL with mAb 9.3, and then removing the CD28⁺ cells with goat anti-mouse coated magnetic beads (Advanced Magnetics Institute, Cambridge, MA). Northern (RNA) blot analysis was done as described

- 30 -

by June, C.H. *et al.*, *Mol. Cell. Biol.* 7:4472-4481 (1987). The IL-3 probe was a 1.0 kb Xho I cDNA fragment.

- To determine if stimulation of the TCR/CD3 pathway of T cell activation induced IL-3 gene expression, CD28⁺ T cells were stimulated with maximal amounts of plastic immobilized anti-CD3 mAb in the presence or absence of 9.3 mAb 1 μ g/ml for 1 to 36 h. As shown in Table 4, anti-CD28 resulted in a 3 to 5-fold augmentation of IL-3 mRNA expression over that induced by anti-CD3 alone. CD28 did not change the kinetics of IL-3 gene expression, which was at peak levels at 6 h after anti-CD3 or after anti-CD3 + anti-CD28 treatment.
- Further experiments showed that IL-3 gene expression was restricted to the CD28⁺ subset of T cells, as determined by Northern analysis (Table 4). The stability of IL-3 mRNA was also determined. T cells were treated for 3 h with anti-CD3 or anti-CD3 plus anti-CD28 mAb to induce IL-3 mRNA expression. At 3 h, actinomycin D was added to the culture to inhibit further RNA synthesis.
- Total cellular RNA was isolated, and the remaining IL-3 mRNA determined by Northern analysis. The half-life of IL-3 mRNA from anti-CD3 plus anti-CD28 treated cells was at least 8-fold longer than the IL-3 mRNA from anti-CD3 treated cells (Table 4). Thus, as would be expected from the previously described results of anti-CD28 on other lymphokines, it can be concluded that the effect of anti-CD28 on IL-3 gene induction can, in large part, be explained by the ability of anti-CD28 to stabilize the IL-3 messenger RNA.

- 31 -

TABLE 4

CONDITION		IL-3 GENE EXPRESSION (arbitrary densitometry units)
EXPERIMENT #1		
5	CD28 ⁺ T cells, 6 h, anti-CD3	1
	CD28 ⁺ T cells, 6 h, anti-CD3 + anti- CD28	3 - 5
EXPERIMENT #2		
10	CD28 ⁺ T cells, 8 h, PMA 3ng/ml + Ionomycin 0.4µg/ml	> 10
	CD28 ⁻ T cells, 8 h, PMA 3ng/ml + Ionomycin 0.4 µg/ml	< 1
EXPERIMENT #3		
15	CD28 ⁺ T cells, anti CD3, then actinomycin D 90 m	1
	CD28 ⁺ T cells, anti-CD3 + anti-CD28, then actinomycin D 90 m	8

SPECIFIC EXAMPLE IV

Comparison of CD28 Stimulation to Stimulation of Other T Cell Surface

20 Molecules.

CD28⁺ T cells were cultured at approximately 1×10^5 cells/well in RPMI media containing 5% heat-inactivated fetal calf serum (FCS), PHA 10 μ g/ml, PMA 3 ng/ml, ionomycin at 100 ng/ml, anti-CD28 mAb 9.3 at 100 ng/ml, or mAb 9.4 specific for CD45 at 1 μ g/ml or mAb 9.6 specific for CD2 at 1 μ g/ml, or
 25 immobilized mAb G19-4 specific for CD3 at 200 ng/well.

CD28⁺ T cells were cultured in quadruplicate samples in flat-bottomed 96-well microtiter plates in RPMI media containing 5% heat-inactivated fetal calf serum. Equal aliquots of cells were cultured for 18 h and then pulsed for 6 h with 1 μ Ci/well of ³H-uridine, or for 72 h and then pulsed for 6 h with 1 μ Ci/well
 30 of ³H-thymidine. The means and standard deviations (in cpm) were determined by liquid scintillation counting after cells were collected on glass fiber filters.

All cultures containing cells immobilized to plastic by anti-CD3 monoclonal antibodies were visually inspected to ensure complete cell

- 32 -

harvesting. The failure of cells in these cultures to proliferate in response to PHA is the result of rigorous depletion of accessory cells, *in vivo* activated T cells, B cells, and CD11⁺ (CD28⁺) T cells by negative immunoabsorption as described in Specific Example II above. In each experiment, cells were stained
5 with fluorescein-conjugated anti-CD2 mAb OKT11 and fluorescein-conjugated anti-CD28 mAb 9.3 and were shown to be over 99% and over 98% surface positive, respectively.

A representative experiment is illustrated in Figures 1 and 2. As shown in Figures 1 and 2, anti-CD28 by itself had no significant effect on uridine or
10 thymidine incorporation, nor did it serve to augment proliferation induced either by immobilized anti-CD3 mAb G19-4 or chemically-induced T cell proliferation involving phorbol myristate acetate (PMA) and ionomycin (Iono). However, as shown in Figure 2, anti-CD28 did significantly increase the uridine incorporation of both sets of cells. In contrast, other monoclonal antibodies including anti-
15 CD2 mAb OKT11 and anti-CD45 mAb 9.4 had no significant effect on uridine incorporation of anti-CD3 stimulated cells. This was not due to lack of effect of these antibodies on the cells, since anti-CD2 monoclonal antibodies significantly augmented the proliferation of anti-CD3 stimulated cells. In separate
20 experiments, the binding of isotype-matched mAbs to other T cell surface antigens (CD4, CD6, CD7 or CD8) failed to mimic the effects observed with anti-CD28.

These data serve to confirm that the stimulation of activated T cells by CD28 has a unique phenotype which appears to directly enhance the rate of incorporation of a radioactive marker into the steady state RNA of T cells
25 without directly enhancing T cell proliferation.

SPECIFIC EXAMPLE V

Increased Cellular Production of Human T_HCD28 Lymphokines by CD28 Stimulation *Ex Vivo*.

Based on evidence from the *in vitro* systems it appeared that CD28 did
30 not have a significant effect on cellular production of lymphokines unless they had undergone prior antigen activation or its equivalent. However, CD28 binding by the 9.3 mAb significantly enhanced the ability of anti-TCR/CD3 activated T cells to sustain production of human T_H1-type lymphokines. To test

- 33 -

this effect in a physiologic setting, the activation of T lymphocytes in an *ex vivo* whole blood model was studied.

50-100 ml of venous blood was obtained by standard aseptic procedures from normal volunteers after obtaining informed consent. The blood was
 5 heparinized with 25 U/ml of preservative-free heparin (Spectrum, Gardenia, CA) to prevent clotting. Individual 10 ml aliquots were then placed on a rocking platform in a 15 ml polypropylene tube to maintain flow and aeration of the sample.

To assay for the effectiveness of CD28 stimulation on the induction of
 10 lymphokine gene expression, the production of TNF- α molecule was chosen as a model because of the extremely short half-life (approximately 15 minutes) of the protein in whole blood. 10 ml of whole blood isolated as described above was incubated with soluble anti-CD3 mAb G19-4 at a concentration of 1 μ g/ml or anti-CD28 mAb 9.3 at a concentration of 1 μ g/ml or a combination of the two
 15 antibodies. The plasma was assayed for TNF- α as described in Specific Example III at one and four h. An example of one such experiment is shown in Table 5, which illustrates the significant increase in sustained production of TNF- α by maximal stimulation of CD3 and costimulation of CD28.

TABLE 5

20	STIMULUS	TNF- α (pg/ml)		
		0 h	1 h	4 h
	anti-CD3	4.5*	65.0	2.1
	anti-CD28	4.5*	1.6	3.3
25	anti-CD3 + anti-CD28	4.5*	35.0	75.0

* value determined prior to addition of monoclonal antibody to aliquots of the venous sample

SPECIFIC EXAMPLE VI

30 Resistance of CD28-Induced T Cell Proliferation to Cyclosporine.

The protocol used and results described herein are described in detail in June, C.H. *et al.*, *Mol. Cell. Biol.*, 7: 4472-4481 (1987), herein incorporated by reference.

- 34 -

T cells, enriched by nylon wool filtration as described by Julius, *et al.*, *Euro. J. Immunol.*, 3:645-649 (1973), were cultured at approximately 5×10^4 /well in the presence of stimulators in the following combinations: anti-CD28 mAb 9.3 (100ng/ml) and PMA 1 (ng/ml); or immobilized anti-CD3 mAb G19-4 (200ng/well); or PMA (100ng/ml). The above combinations also included fourfold titrations (from 25ng/ml to $1.6 \mu\text{g/ml}$) of cyclosporine (CSP) (Sandoz, Hanover, NJ) dissolved in ethanol-Tween 80 as described by Wiesinger, *et al.*, *Immunobiol.*, 156:454-463 (1979).

^3H -thymidine incorporation was measured on day 3 of culture and the results representative of eight independent experiments are depicted in Figure 3. The arithmetic mean ($91,850 \pm 1300$ (mean \pm SD)) CD28-induced T cell proliferation exhibits nearly complete cyclosporine resistance when accompanied by the administration of PMA. Table 6 below illustrates the effects of cyclosporine on CD3-induced proliferation of CD28⁺ T cells cultured at approximately 5×10^4 cells/well in flat-bottomed 96-well microtiter plates (CoStar, Cambridge, MA) under the following conditions: immobilized Mab G19-4; or immobilized mAb G19-4 and mAb 9.30 100ng/ml; or immobilized mAb G19-4 and PMA 1ng/ml; or Mab 9.3 100ng/ml and PMA 1ng/ml. Cyclosporine was prepared as above and included in the cultures at 0, 0.2, 0.4, 0.8, 1.2 $\mu\text{g/ml}$.

^3H -thymidine incorporation was determined on day 3 of culture as above. The percent inhibition of proliferation was calculated between CD28⁺ T cells cultured in medium only or in cyclosporine at 1.2 $\mu\text{g/ml}$. CD28⁺ T cells cultured in the absence of cyclosporine were given cyclosporine diluent. ^3H -thymidine incorporation of cells cultured in medium, or PMA, or monoclonal antibody 9.3 only were less than 150 cpm. As shown in Table 6, costimulation of CD3 and CD28 resulted in a marked increase in the resistance of T cell proliferation to cyclosporine and the stimulation of CD28 in the presence of PMA resulted in a complete absence of cyclosporine suppression of T cell proliferation. As shown in Table 7, stimulation of CD28 together with immobilized anti-CD3 also resulted in resistance to suppression of T cell proliferation by the immunosuppressant dexamethasone.

TABLE 6

Effects of CD28 Stimulation on Cyclosporine Resistance on T Cell Proliferation

Mean [³ H]thymidine incorporation (kcpm) \pm 1 SD at Cyclosporine Conc (ug/mL)						
STIMULUS	0	0.2	0.4	0.8	1.2	% INHIBIT
CD3 mAb G19-4	77 \pm 26	61 \pm 6.8	52 \pm 4.4	10 \pm 3.4	8.2 \pm 1.2	90
CD3 + CD28 mAb 9.3	123 \pm 18	86 \pm 2.3	63 \pm 4.4	44 \pm 6.4	43 \pm 5.2	65
CD3 + PMA	145 \pm 12	132 \pm 2.8	123 \pm 6.4	55 \pm 3.6	56 \pm 6.4	62
CD28 mAb 9.3 + PMA	111 \pm 12	97 \pm 5.6	107 \pm 12	99 \pm 14	112 \pm 2.4	< 0

- 36 -

TABLE 7

Effects of CD28 Stimulation on Dexamethasone Resistance on T cell Proliferation

5

³ HTdR + Dexamethasone (Nm)			
STIMULUS	0	25	%INHIBIT
CD3 mAb G194	14,700	770	97
CD3 mAb + IL-2	21,700	1,900	93
CD28 mAb + PMA	181,600	197,700	<0
PMA	5,000	1,400	72

SPECIFIC EXAMPLE VII

Human T_HCD28 Lymphokine Secretion in the Presence of Cyclosporine.

As described in Specific Example III, CD28⁺ T cells were cultured in the presence of various stimulators. Culture supernatants were harvested at 24 h and serial dilutions assayed for IL-2, TNF- α /LT, IFN- γ , and GM-CSF as previously described. Separate aliquots of cells were recovered 48 h after stimulation and assayed for the percentage of cells in late stages of the cell cycle (S + G₂ + M).

When cyclosporine at 0.6 μ g/ml was included in the test protocol, as shown in Table 8 (which also incorporates the data of Specific Example III for comparison), CD28⁺ T cells were found to secrete the T_HCD28 lymphokines in the presence of cyclosporine in cultures stimulated with mAb 9.3 and PMA; or immobilized mAb G19-4 and mAb 9.3; or PMA and ionomycin and mAb 9.3. T_HCD28 lymphokine production induced by immobilized mAb G19-4; or by PMA with ionomycin was, however, completely suppressed in the presence of cyclosporine.

- 37 -

TABLE 8

	STIMULUS	IL-2 (U/ml)	TNF- α /LT (U/ml)	IFN- γ (U/ml)	GM-CSF (U/ml)	S+G ₂ +M (%)
	Medium	<2	0	0	0	4.6
	PMA	<2	0	0	NT	5.5
5	anti-CD28	<2	5	0	0	6.5
	anti-CD28 + PMA	435	300	24	150	48.9
	anti-CD28 + PMA + CSP	192	200	12	NT	49.3
	anti-CD3 ⁱ	36	50	24	120	39.7
10	anti-CD3 ⁱ + CSP	<2	0	0	NT	14.5
	anti-CD3 ⁱ + anti- CD28	1200	40	74	1050	44.7
	anti-CD3 ⁱ + anti- CD28 + CSP	154	200	9	NT	48.6
15	Ionomycin	<2	0	0	NT	6.6
	Ionomycin + PMA	200	5	37	NT	43.6
	Ionomycin + PMA + CSP	<2	0	0	NT	8.1
20	Ionomycin + PMA + anti-CD28	1640	320	128	NT	43.5
	Ionomycin + PMA + anti-CD28 + CSP	232	120	15	NT	47.6

i = immobilized
 25 NT = not tested

SPECIFIC EXAMPLE VIII

Human T_HCD28 Lymphokine mRNA Expression in the Presence of Cyclosporine.

In order to further examine whether CD28 stimulation led to cyclosporine-
 30 resistant T_HCD28 lymphokine gene expression as well as secretion, the ability of
 cyclosporine to suppress induction of IL-2, TNF- α , LT, IFN- γ , and GM-CSF
 following stimulation by various stimulators was tested. Specifically, CD28⁺ T
 cells were cultured at 2×10^6 /ml in complete RPM1 medium (GIBCO, Grand
 Island, NY) with 5% FCS (MED). Individual aliquots of CD28⁺ T cells were
 35 incubated for 6 h in the presence or absence of 1.0 μ g/ml cyclosporine with

- 38 -

PMA 3ng/ml and anti-CD28 mAb 9.3 (1mg/ml); or with immobilized anti-CD3 mAb G19-4 (1 μ g/well); or with immobilized mAb G19-4 (1 μ g/well) and mAb 9.3 (1ng/ml). CD28⁺ T cells were harvested, total cellular RNA isolated and equalized for ribosomal RNA as previously described by Thompson, *et al.*,

5 *Nature*, 314:363-366 (1985).

Northern blots were prepared and hybridized sequentially with ³²P-labeled, nick-translated gene specific probes as described by June, C.H. *et al.*, *Mol. Cell. Biol.*, 7:4472-4481 (1987). The IL-2 probe was a 1.0 kb Pst I cDNA fragment as described by June, C.H. *et al.*, *Mol. Cell. Biol.*, 7:4472-4481 (1987); the IFN- γ probe was a 1.0 kb Pst I cDNA fragment as described by Young, *et al.*, *J. Immunol.*, 136:4700-4703 (1986). The GM-CSF probe was a 700 base pair EcoR I-Hind III cDNA fragment as described by Wong, *et al.*, *Science*, 228:810-815 (1985); the 4F2 probe was a 1.85 kb EcoR I cDNA fragment as described by Lindsten, *et al.*, *Mol. Cell. Biol.*, 8:3820-3826 (1988); the IL4 probe was a 15 0.9 kb Xho I cDNA fragment as described by Yokota, *et al.*, *PNAS (USA)*, 83:5894-5898 (1986); and the human leukocyte antigen (HLA) probe was a 1.4 kb Pst I fragment from the HLA-B7 gene as described by Lindsten, *et al.*, *Mol. Cell. Biol.*, 8:3820-3826 (1988). TNF- α and LT specific probes were synthesized as gene-specific 30 nucleotide oligomers as described by Steffen, 20 *et al.*, *J. Immunol.*, 140:2621-2624 (1988) and Wang, *et al.*, *Science*, 228:149-154 (1985). Following hybridization, blots were washed and exposed to autoradiography at -70°C. Quantitation of band densities was performed by densitometry as described in Lindsten, *et al.*, *Mol. Cell. Biol.*, 8:3820-3826 (1988).

25 As illustrated by the Northern blot of Figure 4, stimulation by mAb 9.3 with PMA and by mAb 9.3 with mAb G19-4 led to human T_HCD28 lymphokine gene expression that exhibited resistance to cyclosporine. In contrast, stimulation by TCR/CD3 mAb G19-4 alone was completely suppressed in the presence of cyclosporine.

30

SPECIFIC EXAMPLE IX

In Vivo Activation of T Cells by CD28 Stimulation.

A. Monoclonal Antibody 9.3 F(ab')₂.

F(ab')₂ fragments of mAb 9.3 were prepared as described by Ledbetter, J. A. *et al.*, *J. Immunol.*, 135:2331-2336 (1985). Purified and endotoxin-free

- 39 -

F(ab')₂ fragments were injected intravenously at 1 mg/kg of body weight over a 30 minute period into a healthy macaque (*M. nemestrina*) monkey. On days 2 and 7 after injection, 5 ml of blood was drawn and tested.

Peripheral blood lymphocytes from the monkey's blood were isolated by density gradient centrifugation as described in Specific Example II. Proliferation of peripheral blood mononuclear cells in response to PMA (1ng/ml) was tested in the treated monkey and a control animal (no F(ab')₂ fragment treatment) in triplicate as described in Specific Example IV. Proliferation was measured by the uptake of ³H-thymidine during the last 6 h of a three-day experiment and the results shown in Figure 5. Means of triplicate culture are shown, and standard errors of the mean were less than 20% at each point. As shown in Figure 5, *in vivo* stimulation of CD28 by the F(ab')₂ mAb 9.3 fragment increased T cell proliferation for at least 7 days.

B. Monoclonal Antibody 9.3

Two doses of mAb 9.3, 10mg and 0.1mg, were administered intravenously to primates *Macaca mulatta*. The antibody was infused intravenously immediately after baseline (time zero) blood values were obtained. Three animals were evaluated as described below at each dose.

Cell population changes. At the higher dose, immediate effects were monitored over the first 120 m. In Figures 6A and 6B, a representative result is depicted showing the change in the lymphocyte counts over time. The ALC and distribution of CD28⁺ cells are depicted in Figure 6A, while Figure 6B illustrates the absolute numbers of CD4⁺ and CD8⁺ cells. The absolute lymphocyte count (ALC) decreased over the first 60 m and then increased above baseline at 24 h (see Figure 6A). In this case, the number of circulating CD28 positive lymphocytes remained essentially the same, as determined by adding goat anti-mouse phycoerythrin (GAM-PE) only or mAb 9.3 plus GAM-PE. In this same animal, the CD4 and CD8 positive populations were followed and the increase at 24 h was the result of an increase in CD8⁺CD28⁺ cells (see Figure 6B). Animals re-evaluated after 8 days had between 35 to 60% of the CD28⁺ cells coated with antibody. There was no significant change in the circulating lymphocyte counts in primates treated with 0.1mg mAb 9.3.

Cytokine released after *in vitro* stimulation. PBLs isolated at specific time points from primates previously immunized with tetanus toxoid and treated with

- 40 -

10mg mAb 9.3 were cultured *in vitro* to determine the effect of antigenic stimulation on cytokine production. Figure 7A represents the *in vitro* production of TNF while Figure 7B represents the *in vitro* production of IL-6. PBLs stimulated with Concanavalin-a (Con-a) are depicted by Δ . PBLs stimulated with tetanus toxoid (TT) are depicted by \bullet . Unstimulated PBLs are depicted by \circ . As shown in Figures 7A and 7B, cultures of baseline cells did not respond to either Con-a or TT stimulation. However, as shown in Figure 7A, PBLs isolated from animals 6 h after infusion of antibody showed an increase in TNF production. As depicted in Figure 7B, after 24 h, unstimulated cultures produced TNF but not IL-6. TT stimulation of PBLs produced a similar quantity of both TNF and IL-6 as mitogen stimulated cultures. This response was consistent through 72 h.

Monoclonal antibody 9.3 was administered to the primates either as a single day bolus of 10mg (n=3) or as multiple daily injections of 10 mg/d for 5 consecutive days (n=3) and the animals were followed simultaneously. The changes in the peripheral blood cell populations were not dramatic. ALC as previously observed decreased with the first injection but recovered to above baseline if no further injections were administered. However, for animals treated with multiple injections, ALC remained ~25% below normal during the period of mAb administration. ALC in multiple-treated animals did not recover to above normal levels over the 21-day study period. Absolute neutrophil count decreased by ~30% in the 5 day treated group.

Cytokine levels in serum. Serum was analyzed for IL-6, TNF, and IL-1 β . The detection of TNF from the serum preparations was not successful and therefore no results are available at this time. Figures 8A and 8B demonstrate the serum concentration of IL-6 after infusion of mAb 9.3. As shown in Figures 8A and 8B, increased IL-6 levels 24 h after mAb infusion were detected. In animals injected one time, the IL-6 levels increased to a peak on day 4, but a decrease was observed when remeasured on day 8 (see Figure 8A). In comparison, 5 day treated animals (multiple doses) demonstrated continual increase(s) in IL-6 through day 8 (see Figure 8B).

Figures 9A and 9B demonstrate the serum concentration of IL-1 after infusion of mAb 9.3. As shown in Figures 9A and 9B, measurements of IL-1 β in the serum, did not detect any IL-1 until after day 8 in single injected animals

- 41 -

(see Figure 9A) or multiple injected animals (see Figure 9A). The multiple injected animals, however, had increasing levels of IL-1 β at day 21 post-infusion, while single injected animals had decreasing levels at this time.

Cytokine release after *in vitro* stimulation. IL-6 production was not detected in the PBLs of animals on day 3 after *in vitro* stimulation with TT. (Figure 10). This finding contrasted with the previous *in vitro* results. However, an increased production of IL-6 was detected on day 7 and the more significant increase was observed from PBLs isolated from the 5 day treated primates. This increase in production was further observed in culture of PBLs isolated on day 14.

IL-6 Production and Proliferation of PBLs. Figures 10A and 10B illustrate IL-6 production of *in vitro* stimulated PBLs isolated from monkeys. Days 1, 3 and 14 are depicted in Figures 10A and 10B with Δ representing the control and O representing the stimulated PBL response. Figure 10A illustrates the response of a single injected animal and Figure 10B illustrates the response of a multiple injected animal. (Note that the quantity of cells harvested from PBL limited the number of assays performed, resulting in no day zero points and no day zero data.) PBLs were isolated from the different treated groups and evaluated for their proliferative response to Con-A, TT or no stimulus. Historically the TT response was $\sim 5,000$ cpm and the baseline Con-A response was $\sim 35,000$ cpm. The PBL proliferative response to Con-A was reduced by about 80% and gradually recovered over time (not shown). No proliferative response was observed when the PBL were stimulated with TT. This contrasts with the lymphokine production observed in *in vitro* cultures.

25 SPECIFIC EXAMPLE X

Immunoregulation with CTLA-4Ig.

A. *In Vitro*.

The effects of CTLA-4Ig on the primary immune response to alloantigen was initially examined in a one-way mixed lymphocyte culture (MLC) between Lewis rats (RT1^l, responder) and Brown-Norway rats (RT1ⁿ, stimulator). Lymphocytes were isolated from paratracheal and cervical lymph nodes. Cultures were performed in quadruplicate in 96-well round bottomed plates as described in Turka, L.A. *et al.*, *Transplant.*, 47:388-390 (1989). Cultures were harvested after 4 days and 1mCi/well of ³H-thymidine was added for the last 6 h

- 42 -

f culture. In this assay, Brown-Norway stimulator cells were irradiated at 30 Gy to prevent their proliferation, and then added to cultures of Lewis responder lymphocytes. A proliferative response will normally occur in approximately 1-5% of cells as a result of activation through their cell-surface TCR in response to allogeneic MHC as discussed in Marrack, P. *et al.*, *Immunol. Today*, 9:308-315 (1988). Graded concentrations of CTLA-4Ig or an isotype-matched control monoclonal antibody L6 described in Fell, H.P. *et al.*, *J. Bio. Chem.* (in press), was added to the cultures.

Figure 11 represents the effect of CTLA-4Ig on a one-way mixed lymphocyte culture. Spontaneous proliferation is the incorporation of thymidine by Lewis cells in the absence of Brown-Norway stimulators, and is depicted by closed triangles in Figure 11. As shown in Figure 11, CTLA-4Ig was able to block proliferation in a dose dependent fashion with virtually complete inhibition observed at a concentration of 1 mg/ml. (Results are expressed as counts per minute of ^3H -thymidine incorporation \pm standard deviation). Consistent with these results, alloreactive T cell responses can also be inhibited by non-stimulatory Fab fragments of an anti-CD28 monoclonal antibody as shown in Azuma, M. *et al.*, *J. Exp. Med.*, 175:353-360 (1992). Together, these data suggest that in order to mount a proliferative response *in vitro*, alloreactive T cells must be stimulated not only through MHC engagement of the TCR but also require costimulation by B7 engagement of the CD28 receptor.

B. *In Vivo*: Cardiac Allografts.

CTLA-4Ig was next used in a rat model of organ transplantation to ascertain its ability to block alloantigen responses *in vivo*. Recipient animals received a heterotopic cardiac allograft which was anastomosed to vessels in the neck as described in Bolling, S.F. *et al.*, *Transplant.*, 53:283-286 (1992). Grafts were monitored for mechanical function by palpation and for electrophysiologic function by electrocardiogram. Graft rejection was said to occur on the last day of palpable contractile function. As an initial test, animals were treated with daily injections of CTLA-4Ig or an isotype-matched negative control monoclonal antibody L6 for 7 days. CTLA-4Ig was administered at doses of 0.015 mg/day (5 animals), 0.05 mg/day (5 animals), and 0.5 mg/day (8 animals). L6 was given at 0.5 mg/day. Untreated Lewis rats rejected the heterotopic Brown-Norway allografts in 6.8 ± 0.3 days ($n=10$). The allografts

- 43 -

in CTLA-4Ig-treated animals remained functional following completion of drug administration, whereas untreated animals, or animals treated with the L6 control antibody, uniformly rejected their grafts by day 8 ($p < 0.0001$) as shown in Table 9. (p values were calculated by Chi-square analysis).

5

TABLE 9

10

GRAFT SURVIVAL DAY 8		SIGNIFICANCE
Untreated	0/10	$p < 0.0001$
CTLA-4Ig	18/18	$p < 0.0001$
Control Protein	0/5	

15 CTLA-4Ig-treated rats manifested no observable acute or chronic side effects from administration of the protein. No gross anatomic abnormalities were observed in CTLA-4Ig-treated animals at autopsy.

An untreated animal and a CTLA-4Ig-treated animal were sacrificed for histological examination. Cardiac allografts were removed from an untreated animal (shown in Figure 12A) and a CTLA-4Ig-treated animal (0.5 mg/day) (shown in Figure 12B) four days after transplantation. Allografts were fixed in formalin, and tissue sections were stained with hematoxylin-eosin. (Original photography at 200X magnification.) The donor heart removed from the untreated animal showed histological findings of severe acute cellular rejection, including a prominent interstitial mononuclear cell infiltrate with edema formation, myocyte destruction, and infiltration of arteriolar walls. In contrast, the transplanted heart from the CTLA-4Ig-treated animal revealed only a mild lymphoid infiltrate. Frank myocyte necrosis and evidence of arteriolar involvement were absent. The native heart from each animal showed no histological abnormalities.

30 To determine whether CTLA-4Ig therapy established a state of graft tolerance that persisted following drug treatment, animals treated for 7 days with daily injections of CTLA-4Ig were observed without additional therapy until cessation of graft function. Animals received either no treatment, CTLA-4Ig (0.5 mg/day x 7 days), or an isotype-matched control monoclonal antibody, L6

- 44 -

(0.5 mg/day x 7 days). In all cases treatment was initiated at the time of transplantation. Figure 13 is a graph showing allograft survival in the treated and control rats. Graft survival was 18 - 40 days in animals treated with 0.05 mg/day of CTLA-4Ig. Graft survival was assessed daily. This failure to induce permanent engraftment did not appear to be due to inadequate dosing of CTLA-4Ig, as animals treated with a ten-fold higher dose, 0.5 mg/day, showed a similar graft survival curve as depicted in Figure 13, with one animal maintaining long-term graft function (> 50 days). In Figure 13, graft survival is displayed as the last day of graft function. Animals treated with a dose of 0.015 mg/day x 7 days (n=5) had a mean survival of 12.6 ± 2.1 days (n=5). Furthermore, serum CTLA-4Ig trough levels in this group as measured in a quantitative ELISA assay were in excess of 10 $\mu\text{g/ml}$, a concentration which is maximally suppressive *in vitro* (see Figure 11). Histological examination of the allografts from CTLA-4Ig-treated animals whose grafts ceased functioning after 18-43 days displayed typical signs of acute cellular rejection, of the same degree of severity as seen in control animals that had rejected their hearts after 7 days. The animal with continued graft function was sacrificed on day 57, and the allograft from this animal failed to reveal any histological abnormalities.

At the time of sacrifice, lymphocytes from the day 57 "tolerant" animal, and from a CTLA-4Ig-treated animal that rejected the heart at day 33, were tested for their functional responses. These responses were compared with those of lymphocytes from a control (non-transplanted) Lewis rat, and results were normalized as a percentage of the control response. In comparison to control animals, lymphocytes from both the "tolerant" and rejecting animal had equivalent proliferative responses to Con-a, (tolerant, 62.5%; rejecting, 51.1%; $p=0.63$, two-tailed T test) and to cells from a third party ACI rat (RT1^m) (tolerant, 160%; rejecting, 213%; $p=0.58$). However a significant disparity was seen in the response to Brown-Norway cells (tolerant, 34.2%; rejecting, 238%; $p<0.005$), suggesting that T cells from animals with functioning grafts were specifically hyporesponsive to donor MHC antigens. The thymus and spleen from the day 57 "tolerant" animal were similar in size and cell number to the non-transplanted control rat, and flow cytometric analyses of thymus, lymph nodes and spleen revealed similar percentages of both CD4⁺ and CD8⁺ T cells in each animal. Splenocytes adoptively transferred from the day 57 "tolerant"

- 45 -

animal into a native Lewis recipient failed to affect the rate at which that animal rejected a Brown-Norway cardiac allograft. Thus, tolerance did not appear to be maintained by suppressor cells in this animal.

The fact that 4 of 5 allograft recipients treated with high dose CTLA-4Ig (0.5 mg/day x 7 days) rejected their grafts within the study period indicated that blockade of the costimulatory molecule B7 did not consistently induce permanent graft tolerance. One possible explanation was that CTLA-4Ig induced a temporary state of non-responsiveness, and that upon recovery, recipient T cells could effect graft rejection. Alternatively, CTLA-4Ig treatment may have resulted in a state of permanent non-responsiveness in circulating T cells by allowing target antigen recognition without B7-dependent costimulation. Newly matured T cells emerging from the thymus after cessation of CTLA-4Ig treatment could not be tolerized by this mechanism, and could mediate graft rejection as a result of B7-costimulated T cell alloreactivity. To differentiate between these two possibilities, rats were thymectomized 3 days prior to cardiac transplantation, and treated with daily injection of CTLA-4Ig (0.5 mg/day x 7 days) following transplantation. These animals rejected their grafts between days 28 and 33, indicating that allograft recipients were not dependent upon the influx of new T cells to initiate an alloimmune response. Thus, it appears that T cells present during the time of CTLA-4Ig treatment can eventually induce graft rejection. This may represent T cell recovery from a temporary state of non-responsiveness, or may reflect the kinetics of T cell trafficking during the CTLA-4Ig treatment period.

C. Synergistic effects with cyclosporine.

Based on the ability to show that a soluble CD28 receptor homologue, CTLA-4Ig, is capable of suppressing cell-mediated responses *in vitro* and *in vivo*, experimentation was performed to determine whether or not this immunosuppressant has additive or synergistic effect with cyclosporine. A mixed lymphocyte reaction (MLR) with Brown-Norway rat lymph node cells as stimulators and Lewis strain rat lymph node cells as responders was measured by measuring tritiated thymidine incorporation 72 h after cocultivation. The ability of CTLA-4Ig at a concentration of 0.1 μ g/ml and cyclosporine at 30 ng/ml alone or in combination was measured. Figure 14 shows 3 H-thymidine incorporation under various conditions. As can be seen in Figure 14, although

- 46 -

either immunosuppressant led to only a partial reduction in the mixed lymphocyte proliferative response (MLR + CTLA-4Ig and MLR + CSP), the combination of the two (MLR + CSP + CTLA-4Ig) completely blocked the mixed lymphocyte reaction between these MHC-incompatible strains. This effect is greater than what would be expected from two immunosuppressive reagents which have additive effects, suggesting that CTLA-4Ig and cyclosporine block T cell activation by independent mechanisms and have a synergistic effect on T cell activation in response to alloantigens.

SPECIFIC EXAMPLE XI

10 Control of Lymphokine Production by Second Messengers.

A variety of second messengers in the regulation of lymphokine production were examined. In particular, a role for the two primary cell secondary messenger systems, the activation of protein kinase C and elevation in intracellular calcium, were characterized as being central regulators of the transcription of lymphokine genes. In addition, specific tyrosine phosphorylation events were identified that may correlate with the generation of alterations in translation and/or mRNA stability. Further investigations into serine and threonine kinases indicate that they may also have a role in the signal transduction events involved in lymphokine production. In contrast, experiments into the regulation by cGMP showed that this agent has relatively non-specific effects on lymphokine production.

Tyrosine phosphorylation events related to CD28 were further studied as described below.

Protocol

25 **Monoclonal antibodies.** Anti-CD2 mAb G19-4 (IgG1), anti-CD28 mAb 9.3 (IgG2a), anti-CD5 mAb 10.2 (IgG2a), and anti-CD45 mAb 9.4 (IgG2a) were produced, purified and in some cases, biotinylated as described in Ledbetter, J.A. *et al.*, *J. Immunol.*, 135:2331 (1985) and Ledbetter, J.A. *et al.*, *J. Immunol.*, 137:3299 (1986). Anti-B7 mAb 133 (IgM) and dilutions of ascites as described in Freedman, A.S. *et al.*, *J. Immunol.*, 139:3260 (1987), were used. Anti-CD3 mAb OKT3 (IgG2a) was absorbed to goat anti-mouse IgG covalently linked to microspheres (KPL, Gaithersburg, MD), by incubation of a $1/10^5$ dilution of pooled ascites with 10^7 beads/ml in HBSS at room temperature, followed by extensive washing.

- 47 -

Cells. The CD28⁺ subset of T cells was isolated from peripheral blood T lymphocytes by negative selection using immunoabsorption with goat anti-mouse Ig-coated magnetic particles as previously described in June, C.H. *et al.*, *Mol. Cell. Biol.*, 7:4472 (1987). This resulted in a population of resting T cells that was >99% CD3⁺ and that did not contain CD2⁺/CD3⁻ cells such as NK cells. The Jurkat T leukemia cell line E6-1 was a gift from Dr. A. Weiss and maintained in complete media, i.e. RPMI 1640 containing 2 mM L-glutamine, 50 µg/ml gentamycin, and 10% FCS (HyClone Laboratories, Logan, UT). In some instances, T cells or Jurkat cells were cultured in complete media, or in complete media with 5 ng/ml PMA (Sigma Chemical Co., St. Louis, MO) or OKT3 beads (± 5 beads/cell) before experiments. The Jurkat J32 cell line (CD2⁺, CD3⁻, CD28⁺) has been described in Makni, H. *et al.*, *J. Immunol.*, 146:2522 (1991). J32 variants (CD2⁺, CD3⁻, CD28⁺) were derived by γ irradiation-induced mutagenesis and immunoselection (see Makni, *supra* (1991)); one such cloned mutant, J32-72.4 is stable in culture. The surface receptor expression of these cells was quantitated by indirect immunofluorescence and analyzed by flow cytometry. The mean log fluorescence intensity for each sample was determined and was converted into linear relative fluorescence units (ΔFL) by the formula $\Delta FL = 10^{(E-C)/D}$, where *E* is the mean log fluorescence intensity of the experimental antibody sample, *C* is the mean log fluorescence intensity of the control antibody sample, *D* is 50 channels/decade. For the TCR/CD3 and CD28 receptors, ΔFL of the J32 cells was 27.0 and 57.0, and for the J32-74.2 cells 1.1 and 40.7. Northern blot analysis of J32-72.4 revealed no detectable TCR-β mRNA, while the expression of the TCR-α, CD3-γ, δ, and ε and TCR ζ mRNA was similar to that of the parental J32 cells (unpublished data).

B7 transfection of CHO cells. CHO cells were transfected with B7 cDNA as previously described in Gimmi, C.D. *et al.*, *PNAS (USA)*, 88:6575 (1991). These cells have previously been shown to stimulate lymphocyte proliferation and lymphokine secretion in a manner that mimics CD28 mAb-induced T cell activation. See Linsley, P.S. *et al.*, *J. Exp. Med.*, 173:721 (1991) and Gimmi, *supra* (1991). Transfected CHO cells showing no B7 expression were recloned and are referred to as CHO-B7⁻. CHO cells were detached from tissue culture plates by incubation in PBS with 0.5 mM EDTA for 30 m and fixed in 0.4%

- 48 -

paraformaldehyde as described in Gimmi, *supra* (1991). Fixed CHO-B7 cells were used as control cells.

Immunoblot analysis of protein tyrosine phosphorylation. Details of the immunoblot assay with anti-phosphotyrosine antibodies has been described in Hsi, E.D. *et al.*, *J. Biol. Chem.*, 264:10836 (1989) and June, C.H. *et al.*, *J. Immunol.*, 144:1591 (1990). Cells were suspended at $5 - 10 \times 10^7$ cells/ml in reaction media, i.e., HBSS containing 0.8% FCS and 20 Mm HEPES at 37°C at time - 3 m and stimulated at time 0 m. mAbs were used at 10 µg/ml final concentration. For crosslinking, biotinylated mAbs were incubated with cells for 5 - 8 m at room temperature, the cells prewarmed at time 3 min and stimulated with avidin (Sigma Chemical Co.) at a final concentration of 40 µg/ml at time 0. Stimulation was terminated by the addition of ice-cold 10x lysis buffer, yielding a final concentration of 0.5% Triton X-100. See June, *J. Immunol.*, *supra* (1990). After lysis at 4°C, nuclei were pelleted and postnuclear supernatants were subject to SDS-PAGE on a 7.5% gel, transferred to polyvinylidene difluoride microporous membrane (Millipore, Bedford, MA) and the membranes probed with affinity-purified anti-phosphotyrosine antibodies, labeled with ¹²⁵I staphylococcal protein A (ICN, Irvine, CA) and exposed to x-ray film.

Results

Herbimycin A prevents CD28-stimulated IL2 production. Previous studies have shown that three distinct biochemical signals, provided by phorbol esters, calcium ionophore, and ligation of the CD28 receptor with mAb, are required to cause optimal IL-2 secretion (see June, C.H. *et al.*, *J. Immunol.*, 143:153 (1989)). Cells cultured in the presence of PMA, ionomycin, or CD28 mAb alone produced no detectable IL-2 and, as previously reported in June, *J. Immunol.*, (1989) *supra*, and Fraser, J.D. *et al.*, *Science* (Wash., D.C.) 251:313 (1991), stimulation of the CD28 receptor strongly up-regulated IL-2 production of T cells stimulated with immobilized anti-CD2 mAb, PMA, or PMA plus ionomycin. To address the potential role of tyrosine kinases in CD28-triggered signaling, the effect of herbimycin A, an inhibitor of the *src* family protein tyrosine kinases (see Uehara, Y. *et al.*, *Biochem. Biophys. Res. Commun.*, 163:803 (1989)), on the CD28-triggered enhancement of IL-2 production was investigated. T cells were cultured overnight in the absence (depicted as open bars in Figure 15) or presence (depicted as filled bars in Figure 15) of herbimycin A (1µM). The cells

- 49 -

were then cultured for a further 24 h period in the presence of medium-immobilized anti-CD3 mAb (G19-4), PMA (3ng/ml) (P), or PMA plus ionomycin (150 ng/ml) (P+I) in the presence or absence of soluble anti-CD28 mAb 9.3 (1ug/ml). Cell-free supernatant was collected, dialyzed to remove herbimycin A and serial dilutions were analyzed for IL-2 content by bioassay as described in June, *J. Immunol.*, *supra* (1989). Figure 15 shows the effect of herbimycin A on CD28-stimulated IL-2 production. The CD28 mAb mediated enhancement of IL-2 production in response to stimulation with immobilized anti-CD3, or PMA was nearly completely inhibited in the presence of herbimycin A. In contrast, cells cultured in PMA, ionomycin or 9.3 mAb only produced < 10 U/ml of IL-2.

Disruption of the proximal signaling pathway triggered through CD3 could potentially explain the effect of herbimycin on cells stimulated with anti-CD3 and anti-CD28. Consistent with this, CD3-triggered IL-2 production was previously shown to be exquisitely sensitive to herbimycin A. See June, C.H. *et al.*, *PNAS (USA)*, 87:7722 (1990). However, IL-2 production induced with the combination of PMA plus ionomycin or PMA plus CD28 stimulation permits, in principle, the ability to isolate the CD28 signal for testing the effect of herbimycin A. PMA plus anti-CD28-stimulated IL-2 production was sensitive to the effects of herbimycin A while, as previously noted, PMA plus ionomycin-stimulated IL-2 secretion was resistant to the effects of herbimycin A. The combination of PMA plus ionomycin plus anti-CD28-stimulation resulted in more IL-2 secretion than optimal amounts of PMA plus ionomycin, consistent with the previous reports of June, *J. Immunol.*, *supra* (1989); and Fraser, J.D. *et al.*, *Science* (Wash. D.C.) 251:313 (1991). However, in the presence of herbimycin A, PMA plus ionomycin plus CD28-stimulated cells produced approximately equivalent amounts of IL-2 as cells stimulated in the absence of herbimycin with PMA plus ionomycin. Together, the above results suggest that the function of both the TCR and CD28 receptors are sensitive to herbimycin, and further suggest the independent effects of these three reagents on IL-2 gene expression. See June, *J. Immunol.*, *supra* (1989); and Fraser, *supra* (1991).

CD28 receptor crosslinking with mAb induces protein tyrosine phosphorylation in PMA-treated Jurkat cells. Given the above functional results, the potential involvement of protein tyrosine phosphorylation in CD28-mediated signal transduction was investigated by immunoblot analysis of postnuclear

- 50 -

supernatants of whole cell lysates of the T cell leukemia line Jurkat E6-1. Jurkat E-6 cells were cultured for 2 days in the presence or absence of PMA (5 ng/ml). After washing, 10^7 cells in $120 \mu\text{l}$ were stimulated with reaction media (control), anti-CD3 Mab (G19-4), anti-CD28 mAb (9.3), or crosslinked anti-CD28 mAb (9.3) (final concentration, $10 \mu\text{g/ml}$). For crosslinking, biotinylated mAb was added at time 10 m, followed by avidin ($40 \mu\text{g/ml}$) at time zero. After 2 m, the reaction was terminated with ice-cold lysis buffer and postnuclear supernatants were resolved by SDS-PAGE electrophoresis, transferred to immobilon, and immunoblotted with antiphosphotyrosine, followed by ^{125}I -protein A and autoradiography.

In a previous report by Ledbetter, J.A. *et al.*, *Blood*, 75:1531 (1990), increased tyrosine phosphorylation could not be detected in resting T cells after crosslinking the CD28 receptor. Consistent with that report, no changes in tyrosine phosphorylation were detected in unstimulated Jurkat cells after the binding of bivalent or crosslinked CD28 mAb. Previous studies have shown that CD28 stimulation alone does not result in lymphokine production in Jurkat cells or induce proliferation of primary T cells. See Weiss, A. *et al.*, *J. Immunol.*, 137:819 (1986); Martin, P.J. *et al.*, *J. Immunol.*, 136:3282 (1986); and Hara, T. *et al.*, *J. Exp. Med.*, 161:1513 (1985). Engagement of CD28 by CD28 mAbs or by B7, the natural CD28 ligand, delivers a costimulatory signal provided T cells are stimulated with PMA or with TCR/CD3 mAbs. See June, C.H. *et al.*, *Immunol. Today*, 11:211 (1990); Koulova, L. *et al.*, *J. Exp. Med.*, 173:759 (1991); Linsley, P.S. *et al.*, *J. Exp. Med.*, 173:721 (1991); and Gimmi, C.D. *et al.*, *PNAS (USA)* 88:6575 (1991). It thus appeared that CD28-induced protein tyrosine phosphorylation might only occur in the context of a costimulatory signal.

To test this hypothesis, Jurkat cells were cultured in PMA and then stimulated with anti-CD28 mAb as previously described. In the PMA-stimulated cells, crosslinking of CD28 for 2 m induced phosphotyrosine on substrates migrating with approximate molecular masses of 47, 62, 75, 82, 100, 110, and 145kD. Bivalent CD28 mAb induced tyrosine phosphorylation, but to a lesser magnitude. In agreement with June, C.H. *et al.*, *J. Immunol.*, 144:1591 (1990), CD3 triggering of Jurkat cells induced tyrosine phosphorylation of phosphoprotein (pp) 56, pp65, pp75, pp100, pp110, and pp145 in resting

- 51 -

Jurkat cells and in PMA-treated Jurkat cells. Of particular interest were pp75 and pp100, which were consistently phosphorylated by CD28 stimulation under all conditions tested.

CD28 receptor crosslinking with Mab induces protein tyrosine

- 5 ***phosphorylation in normal T cells.*** Similar experiments with highly purified peripheral blood T cells from normal human donors were performed in order to determine if CD28 could increase tyrosine phosphorylation in nontransformed cells. Peripheral blood CD28⁺ T cells were cultured in PMA (5 ng/ml) for 6 h. After washing, 10⁷ cells were stimulated for 2 m with media (control), anti-CD3
10 mAb (G19.4), anti-CD28 mAb (9.3), crosslinked anti-CD28 mAb (9.3), or crosslinked anti-CD5 mAb (10.2). Cells were lysed and protein tyrosine phosphorylation was determined as previously described. Crosslinking of CD28 on PMA-treated cells induced the appearance of tyrosine phosphorylated substrates that migrated at 45, 75, and 100kD. Again, pp75 and pp100 were
15 most prominent and consistently reproduced.

- The effects of CD28 stimulation observed after 24 - 48 h of PMA stimulation were more pronounced than those seen after 6 h. Ligation of CD28 by mAb on resting T cells caused the appearance of weakly detected tyrosine phosphorylation. The induction of increased responsiveness to anti-CD28 mAb
20 stimulation by PMA is slow in that 4 - 6 h of PMA treatment are required to consistently observe CD28-induced tyrosine phosphorylation. Experiments with cycloheximide indicate that new protein synthesis is required for cells to become responsive to CD28. The specificity of the CD28-induced tyrosine phosphorylation was investigated by crosslinking CD5 with an isotype-matched
25 mAb. Increased tyrosine phosphorylation on the 75kD substrate was occasionally induced by CD5 crosslinking. In contrast, CD5 never induced tyrosine phosphorylation on pp100. Similarly, crosslinking of the MHC class I receptor also did not induce tyrosine phosphorylation of this substrate.

CD28 receptor crosslinking induces protein tyrosine phosphorylation in

- 30 ***CD3-treated normal T cells.*** The above experiments suggested that the CD28 receptor is relatively inactive in quiescent cells, and becomes responsive consequent to protein kinase C activation. To determine whether TCR stimulation could also prime cells for the CD28 signal, T cells were cultured overnight in medium or in the presence of anti-CD3-coated beads. The cells

- 52 -

were recovered, and 8×10^6 cells were stimulated with crosslinked anti-CD28 mAb for 0 - 5 m, the cells lysed, and protein tyrosine phosphorylation determined as previously described. Crosslinked CD28 mAb induced low level tyrosine phosphorylation on multiple substrates in resting T cells that peaked 2 - 5 m after CD28 stimulation. In contrast, CD28 mAb induced marked tyrosine phosphorylation in CD3-primed cells that was maximal within 1 m. Thus, costimulation of T cells with anti-CD3 augmented CD28-induced tyrosine phosphorylation as manifested by an increased magnitude of response and an accelerated kinetics of response. This induction of responsiveness to CD28 did not require DNA synthesis, as separate studies have shown that the T cell blasts used for these studies were in the late G₁ phase of the cell cycle.

CD28 receptor-B7/BB1 receptor interaction induces specific tyrosine phosphorylation in T cells. The above results indicate that CD28 mAb can increase tyrosine phosphorylation in a variety of substrates on preactivated T cells. Previous studies have indicated that CD28 appears to deliver two biochemically distinct signals, depending on the degree of crosslinking. See Ledbetter, J.A. *et al.*, *Blood*, 75:1531 (1990). The unique functional properties of CD28 mAb observed after stimulation of T cells do not require highly crosslinked CD28 mAb and are obtained using intact or F(ab')₂ CD28 mAb. As discussed in Hara, T. *et al.*, *J. Exp. Med.*, 161:1513 (1985), studies have shown that CHO cells expressing the CD28 ligand mimic the functional effects of CD28 mAb. See Linsley, *supra* (1991), and Gimmi, *supra* (1991). These cells presumably represent a more physiologic means to study CD28 receptor-mediated signal transduction. CHO-B7⁺ cells were incubated with PMA-treated T cells at a CHO/T cell ratio of 1:10 for 5 - 30 m. B7-transfected CHO cells not expressing B7 on the cell surface (CHO-B7⁻ cells) were used as controls. Before the stimulation, CHO cells were fixed with paraformaldehyde to decrease phosphotyrosine background. Previous studies have indicated that this treatment leaves intact B7-CD28 interaction and the ensuing functional effects. See Gimmi, *supra* (1991). For the time zero point, lysis buffer was added to the T cells first, immediately followed by addition of CHO cells to the mixture. CHO-B7⁺ cells induced specific tyrosine phosphorylation that was detected primarily on a substrate that migrated at 100kD. The CHO-B7-induced tyrosine phosphorylation was detectable within 5 m of stimulation and remained elevated

- 53 -

at plateau levels for at least 30 m. CHO-B7-induced tyrosine phosphorylation was evident at a variety of CHO-T cell ratios, and has been consistently observed for only the 100kD substrate. CHO-B7 cells did not induce tyrosine phosphorylation of pp100. The B7-induced tyrosine phosphorylation was
5 dependent upon CD28-B7 interaction as preincubation of CHO cells with anti-B7 mAb prevented CHO-B7 induced pp100 tyrosine phosphorylation. B7-CHO cells induced a slight increase in pp100 tyrosine phosphorylation in some experiments; however, this was not consistently observed.

In other experiments, alloantigen-induced T cell blasts were tested for
10 CD28-induced tyrosine phosphorylation. T cells were culture for 8 days with allogeneic irradiated cells and then stimulated with CD28 mAb. Tyrosine phosphorylation that was most pronounced on the 74 and 100kD substrates was observed. Thus, CD28 stimulation of T cells preactivated with alloantigen, CD3 mAb, or PMA can induce tyrosine phosphorylation on a limited number of
15 substrates that is early in onset and brief in duration.

CD28-induced tyrosine phosphorylation prevented by CD45 and by herbimycin. Given that protein tyrosine kinase inhibitor herbimycin A could efficiently inhibit CD28-induced IL-2 secretion, this inhibitor was tested for effects on CD28-induced tyrosine phosphorylation. T cells were treated
20 overnight with PMA (5 ng/ml) in the presence of the indicated concentration of herbimycin A or in control medium. The cells were collected, washed, and 8×10^6 cells were stimulated with media or with crosslinked anti-CD28 mAb for 2 m. Detergent-soluble proteins were processed as previously described. Tyrosine phosphorylation induced by anti-CD28 mAb was nearly completely
25 prevented in herbimycin-treated cells under conditions that specifically inhibit CD28-induced IL-2 production.

The brief temporal course of CD28 mAb-induced tyrosine phosphorylation suggested regulation by a phosphatase. To address the effects of phosphatases on CD28-mediated signal transduction, T cells were cultured overnight with
30 PMA (5 ng/ml). 10^7 cells were incubated for 10 m with media (control), biotinylated anti-CD45 mAb (9.4), anti-CD28 mAb (9.3), or both. Monoclonal antibodies were crosslinked with avidin at time 0. The reaction was terminated after 2 m. Immunoblot analysis with antiphosphotyrosine antibodies of detergent-soluble proteins was performed as previously described. CD28

- 54 -

crosslinking induced tyrosine phosphorylation on pp75 and pp100 that was completely prevented by CD45. Consistent with previous results described in Samelson, L.E. *et al.*, *J. Immunol.*, 145:2448 (1990), crosslinking of CD45 alone caused increased tyrosine phosphorylation of a 120-135kD substrate; this effect is also seen in CD28 plus CD45-treated cells. Thus, the above studies indicate that CD28-induced tyrosine phosphorylation is sensitive to an inhibitor of *src* family protein tyrosine kinases, and furthermore, that the CD45 protein tyrosine phosphatase can prevent CD28-induced protein tyrosine phosphorylation.

- 10 ***CTLA-4 expression predicts expression of IL-2 following CD28 pathway activation.*** Purified resting T cells were stimulated with immobilized anti-CD2 Ab, anti-CD3 + mAb 9.3, PMA + ionomycin, PMA + mAb 9.3 and PMA + ionomycin + mAb 9.3 in the presence or absence of the protein-tyrosine kinase inhibitor herbimycin for 8 h. Duplicate Northern blots were hybridized to CTLA-
15 4, CD28, IL-2 or HLA specific probes. Expression of CD28, CTLA-4 and IL-2 was then analyzed by Northern blot. IL-2 expression correlated well with CTLA-4 expression following CD28 pathway activation. CTLA-4 and IL-2 expression were also suppressed to a similar degree with herbimycin while CD28
20 expression remained unchanged. This suggests that the suppressive effects of protein-tyrosine kinase inhibitors on CD28 pathway activation may be mediated through suppression of CTLA-4 expression.

SPECIFIC EXAMPLE XII

Induction of MHC Independent T Cell Proliferation by CD28 and Staphylococcus Enterotoxins.

25

Protocol

- Isolation of T cells.*** Peripheral blood was drawn from normal human volunteers. The mononuclear cell fraction was obtained by density gradient centrifugation through a Ficoll-Hypaque (Pharmacia) cushion. This fraction was used in experiments utilizing peripheral blood mononuclear cells (PBMCs).
30 Purified resting T cells were obtained by incubating the mononuclear cells with an antibody cocktail directed against B cells, monocytes and activated T cells. The antibody coated cells were then removed by incubation with goat anti-mouse immunoglobulin-coated magnetic beads (Advanced Magnetix Inc.)

- 55 -

as previously described in June, *supra* (1987). This method has routinely yielded a population > 99% CD2⁺ by flow cytometry.

Proliferation assays. Proliferation was measured by culturing 5×10^5 purified T cells or PBMCs in each well of a 96 well microtiter plate. The final culture volume was 200 μ l of RPMI 1640 (Gibco) supplemented with 10% FCS, penicillin (100 U/ml), streptomycin (100 μ g/ml) and 2 Mm L-glutamine. Staphylococcal Enterotoxin A (SEA), Staphylococcal Enterotoxin B (SEB) (Toxin Technologies) and cyclosporine A (Sandoz) were added in the indicated doses at the initiation of the culture. Anti-CD28 monoclonal antibody (mAb 9.3, gift from J. Ledbetter) and anti-HLA-DR monoclonal antibody (mAb L243, gift from J. Ledbetter) were added at the start of the culture period. Tritiated thymidine (³H-TdR, ICN) was included at a concentration of 1 μ Ci per well for the final 8 h of the culture. The cells were harvested onto glass microfiber filter strips (Whatman) after 72 h using a PHD cell harvester (Cambridge Technologies) and counted on a liquid scintillation counter (LKB). All values are expressed as the mean cpm \pm standard deviation of triplicate or quadruplicate cultures.

Flow Cytometry. One ml cultures of T cells were incubated with media alone, SEA (100 ng/ml) or SEB (1 μ g/ml), SEA or SEB plus anti-CD28 antibody (1 μ g/ml) or PMA (3 ng/ml) plus anti-CD28 antibody (1 μ g/ml) at 37°C for 72 h. Aliquots of each sample were stained with acridine orange (Polysciences) for cell cycle analysis as described in Darzynkiewicz, Z., *Meth. Cell. Biol.*, 33:285 (1990), FITC conjugated anti-IL2 receptor antibody (Coulter), FITC conjugated anti-HLA-DR antibody (Becton-Dickinson) or an isotype-matched irrelevant antibody (Becton-Dickinson). Each sample was analyzed on a FACScan flow cytometer (Becton-Dickinson).

Results

CD28 provides costimulatory activity for superantigen-activated purified T cells. Highly purified T cells were cultured with graded concentrations of either SEA (0.1 ng/ml to 1.0 μ g/ml) or SEB (.01 μ g/ml to 100 μ g/ml). Replicate cultures were prepared in which a stimulatory antibody to CD28 was added. The cultures were pulsed with ³H-TdR for the final 8 h of a 72 h culture and incorporated thymidine determined by liquid scintillation counting as described above. Each condition was performed in quadruplicate. Treatment with SEB alone failed to induce thymidine incorporation above control cultures. However,

- 56 -

the addition of anti-CD28 antibody resulted in significant proliferation to graded doses of SEB. Treatment with anti-CD28 antibody alone had no effect. The lack of accessory cells was verified by an absence of proliferation to PHA. Identical results were obtained using SEA.

- 5 ***Stimulation with SEA or SEB leads to cell cycle entry.*** Since CD28 stimulation alone does not induce T cell cycle entry, the observation that CD28 provided costimulatory activity for T cells treated with either SEA or SEB suggested that these enterotoxins could induce cell cycle entry in purified T cells. In order to examine this, purified T cell cultures were stimulated with SEB
- 10 (1 $\mu\text{g/ml}$) alone or with SEB (1 $\mu\text{g/ml}$) plus anti-CD28 monoclonal antibody (1 $\mu\text{g/ml}$) for 48 h and stained with acridine orange for cell cycle analysis. Unstimulated cells were run simultaneously in order to determine the G_0/G_1 interface. Those with an increased RNA content but unchanged DNA content were considered G_1 phase cells. Cells with increases in both RNA and DNA
- 15 content were considered in S, G_2 or M phases. Concomitantly, aliquots were stained with FITC-conjugated anti-IL-2 receptor antibody. Treatment with enterotoxin alone for 48 h resulted in progression of greater than 10% of the T cells from G_0 to G_1 as determined by an increase in RNA staining with no increase in DNA content. Similarly, enterotoxin alone induced IL-2 receptor
- 20 expression in 15% of the cells at 72 h. In contrast, when the anti-CD28 monoclonal antibody was present, a significant proportion of the cells that had left the G_0 stage of the cell cycle were found to have increased their DNA content and thus are in either the S, G_2 or M phases of the cell cycle. These data indicate that stimulation with SEB alone is sufficient to activate the T cell
- 25 but delivers an inadequate signal for complete progression through the cell cycle. Provision of a second signal by simultaneous stimulation of the CD28 pathway allowed the cell to progress to S-phase and proliferate.

- Proliferation of T cells stimulated by SEA and anti-CD28 is resistant to cyclosporine A.*** CD28 has been shown to utilize a signal transduction pathway
- 30 that is resistant to the effects of cyclosporine A (CsA) when the initial signal is provided by PMA, and partially resistant to CsA when cells are initially activated through the T cell receptor. See June, *supra* (1987). In order to further examine the pathways involved in T cell activation by enterotoxin, cyclosporine A (1 $\mu\text{g/ml}$) was included in cultures activated by SEA and SEA plus anti-CD28

- 57 -

antibody (^3H -TdR incorporation determined as described above). As for SEB, SEA alone did not induce thymidine incorporation whereas addition of antibody against CD28 resulted in significant proliferation. Even in the presence of cyclosporine A ($1\text{ }\mu\text{g/ml}$), there was a dose dependent increase in proliferation
5 when cultures were activated by a combination of SEA and anti-CD28 antibody. Control cultures activated with PMA plus anti-CD28 antibody were resistant to cyclosporine A and activation by PMA plus ionomycin was sensitive to cyclosporine A.

Activation by SEB and anti-CD28 is independent of class II MHC.

- 10 Previous work has demonstrated that the staphylococcal enterotoxins are capable of simultaneously binding the TCR and class II MHC molecules on the surface of antigen presenting cells (APCs). See Herrmann, T. *et al.*, *Eur. J. Immunol.*, 19:2171 (1989); and Chintagumpala, M.M. *et al.*, *J. Immunol.*, 147:3876 (1991). The observation that proliferation was not observed unless
15 APCs were present in the culture was interpreted to mean that T cell activation by superantigen is dependent upon class II MHC expression as discussed in Fleischer, B. *et al.*, *J. Exp. Med.*, 167:1697 (1988); Carlsson, R. *et al.*, *J. Immunol.*, 140:2484 (1988); and Herman, A. *et al.*, *J. Exp. Med.*, 172:709 (1990). Our observation that highly purified T cells could be induced to
20 proliferate by simultaneous stimulation with enterotoxin and anti-CD28 antibody suggested that class II MHC may not be absolutely required for superantigen activation of T cells. Alternatively, activated T cells can express class II MHC and thus might provide class II-dependent superantigen presentation to other T cells in trans.
- 25 To examine this possibility, a blocking antibody against HLA-DR, monoclonal antibody L243, was included in cultures of T cells and PBMCs activated by enterotoxin or enterotoxin plus anti-CD28 antibody as shown in Figure 16. Each point is expressed as the mean \pm the standard deviation of triplicate or quadruplicate cultures. No significant proliferation was observed
30 with SEB alone. As shown previously, inclusion of anti-CD28 antibody allows SEB to induce T cell proliferation in a dose-dependent manner. There was no decrease in proliferation when anti-class II antibody was included in the cultures at doses of 1.0 or $10\text{ }\mu\text{g/ml}$ (data for $10\text{ }\mu\text{g/ml}$ not shown). In contrast, as shown in Figure 17, the proliferation of PBMCs isolated from the same donor

- 58 -

and stimulated with enter toxin was significantly inhibited by anti-class II antibody. In addition, HLA-DR expression at 24 and 72 h was examined by purified T cells activated with SEA (0.1 or 1.0 ng/ml) with and without CD28 costimulation. There was no expression of HLA-DR in either condition as
5 determined by flow cytometry. This indicated that the T cell proliferation induced by enterotoxin + anti-CD28 is not dependent on presentation by an MHC class II molecule.

SPECIFIC EXAMPLE XIII

Prevention of Programmed Cell Death.

10 A series of experiments were done to test whether anti-CD28 mAb might prevent cell death in mature T cells. Jurkat leukemia cells are commonly used as an example of mature T cells that mimic physiologic effects found in peripheral blood T cells. For example, Jurkat cells can be induced to secrete IL-2 with anti-CD3 mAb and anti-CD28 mAb, and Jurkat cells can be infected and
15 killed by HIV-1. The Jurkat line JHMI-2.2 was obtained from A. Weiss (UCSF); the muscarinic M₁ receptor subtype has been transfected and is stably expressed in these cells. JHMI-2.2 cells, 0.3×10^6 /well, were added to culture wells in complete medium, or to wells that contained plastic-adsorbed anti-CD3 mAb G19-4, in the presence or absence of 9.3 mAb $10 \mu\text{g/ml}$, or 9.3 mAb
20 alone. Cell death was scored after 1 to 3 days of culture and graded as 0 (none), 1+ (20 to 70% of cells dead), and 2+ (70 to 100% of cells dead), and was determined by visual inspection of the wells, and confirmed by trypan blue permeability. As shown in Table 10, cells in medium continued to grow and remain viable while cells in anti-CD3-treated wells died. In contrast, the cells in
25 wells containing anti-CD3 plus anti-CD28 continued to proliferate. The ability of anti-CD28 to rescue cells from anti-CD3-induced cell death was specific, because carbachol ($30 \mu\text{M}$) (a specific agonist of M₁ receptors that is believed to activate signal transduction in cells via a mechanism distinct from the T cell receptor/CD3 complex), also induced cell death in Jurkat cells. Anti-CD28 did
30 not prevent carbachol-induced cell death.

- 59 -

TABLE 10

CONDITION		CELL DEATH (GRADE 0-2)
EXPERIMENT #1		
5	Medium	0
	anti-CD3	2+
	anti-CD3 + anti-CD28	0
EXPERIMENT #2		
10	Medium	0
	Carbacol	1+
	Carbacol + anti-CD28	2+

SPECIFIC EXAMPLE XIV

Bone Marrow Studies.

Proliferation of T cells after activation of T cells with soluble or immobilized anti-CD3 (OKT3). A series of titration studies were performed using soluble or immobilized OKT3 to activate and induce T cell proliferation. Immobilized OKT3 (2 μ g/ml precoated plates for 1 h at 37°C) and soluble OKT3 (10 ng/ml) consistently induced T cell proliferative responses from E-rosette (E⁺) purified T cells or PBL. PBL or purified T cells were activated by incubation for 1 h to 7 days on immobilized OKT3 or by adding 10 ng/ml of soluble OKT3 at the beginning of culture. In a series of experiments, proliferation after activation of T cells with immobilized OKT3 was comparable to proliferative responses by PBL after activation with soluble OKT3.

Cytotoxicity mediated by anti-CD3 and anti-CD28 triggered PBL. The cytotoxicity of anti-CD3 activated PBL after 7 days of culture in the presence of low doses of IL-2 or anti-CD28 was tested. In this set of experiments, the ability of CD28 to induce increases in lymphokine production to substitute for previously reported immune augmented effects of *in vitro* T cell treatment with IL-2 has been examined. One lytic unit is equivalent to 20% lysis of 5×10^3 target cells per 1×10^6 effector cells as discussed in Press, H.F. *et al.*, *J. Clin.*

- 60 -

Immunol. 1:51-83 (1981). Various targets, including Daudi and K562, were tested. Cytotoxicity results of a representative experiment are shown in Table 11.

TABLE 11

5	CELL GROWTH STIMULUS	Target - Daudi (LU)	Target - K562 (LU)
	anti-CD3 plus IL-2	14.9	10.6
	anti-CD3 plus anti- CD28	12.1	6.1

10

OKT3-induced cytotoxicity in PBL comparable to T cells. In 5 different subjects, PBL were compared with T cells (E⁺) in their ability to kill Daudi, K562, and BSB cells 8 days after being activated with OKT3. These experiments were performed in X-Vivo 10 supplemented with 5% human serum (HS). The mean cytotoxicity in 5 normal subjects using PBL directed at Daudi, K562, and BSB were 15.0, 7.4, and 9.2 LU, respectively. In T cells from the same 5 subjects, the mean cytotoxicity directed at Daudi, K562, and BSB were 14.3, 7.7, and 9.3 LU, respectively.

Cytotoxicity as a function of in vitro time in cell culture. In order to test for optimal cytotoxicity, T cells were cultured with anti-CD3 and IL-2 for 31 days and tested at weekly intervals for cytotoxicity against Daudi and K562. Logarithmic cell growth was maintained during this time, with a 300-fold expansion in cell number. Cytotoxicity was a strong function of culture duration however, with <0.1, 24, 3.5, 0.5, 1.0 LU at 0, 8, 15, 22, and 29 days of culture. Similar results were found when Daudi was the target, with <.01, 23, 5, 2, and 5 LU at 0, 8, 15, 22, and 29 days of culture.

Cytotoxicity induced by soluble or immobilized OKT3. In 7 experiments, cytotoxicity mediated by T cells after activation was compared with soluble or immobilized OKT3. Both methods induced cytotoxicity directed at Daudi and K562. Soluble OKT3 activated T cells mediated a mean cytotoxicity of 27 LU (SD - 18) directed at Daudi and a mean cytotoxicity of 21 LU (SD - 16) directed at K562. Immobilized OKT3 activated T cells mediated a mean cytotoxicity of

- 61 -

22 LU (SD - 16) directed at Daudi and a mean cytotoxicity of 12 LU (SD - 8) directed at K562. These experiments were performed with E⁺ cells in RPMI 1640 supplemented with 10% fetal bovine serum (FBS). Cytotoxicity was assessed 7 to 8 days after triggering with soluble (10 ng/ml) or immobilized (2
5 μ g/ml) OKT3.

Effects of IL-2 concentrations. The dose of IL-2 was titrated after establishing the optimal time of OKT3 activation. In several experiments, the doses of IL-2 were gradually reduced from 6000 IU/ml to 60 IU/ml. Proliferation, as measured by tritiated thymidine incorporation after 3 days of
10 culture, remained constant as IL-2 was titrated in this range. In contrast, cytotoxicity varied with the dose of IL-2, and was maximal at lower doses of IL-2 (lytic units with Daudi targets were 28, 9, 8.5, 9 and 4.8 LU after culture in 30, 150, 300, 600 and 6000 IU/ml of rIL-2). The data show that both
15 proliferative and cytotoxic responses of the anti-CD3 triggered T cells can be obtained and maintained in low doses of IL-2.

Effects of serum and medium on proliferation and cytotoxicity. The ability of HS, FBS, and serum free media were compared for their ability to support growth and maintain cytotoxicity. The data show that proliferation and cytotoxicity directed at Daudi, K562, and BSB rapidly decreased below a serum
20 concentration of 2%. There were no significant differences between X-Vivo 10 and RPMI 1640.

Bone marrow mononuclear cells (BMMNC) as a source of CTC after anti-CD3 and anti-CD28 treatment. To test whether BMMNC might serve as a source of T cells for therapy in patients with malignancies, BMMNC were
25 cultured in X-Vivo 10 medium after OKT3 and IL-2 or anti-CD28 stimulation. Although the BMMNC population initially contained only 25% CD3⁺ cells, proliferative and cytotoxic responses were excellent after 2 weeks of culture. T cells expanded more than 40-fold after CD3 and IL-2 stimulation and cytotoxicity was between 5 and 12 LU at 8 to 15 days of culture when tested
30 against Daudi and K562. These data show that BMMNC obtained from autologous bone marrow harvest from a patient before bone marrow transplantation provide a suitable source of cytotoxic T cells. Table 12 shows that both normal and patient bone marrows provide satisfactory sources of cytotoxicity after CD3 and IL-2 treatment. In 4 experiments using normal bone

- 62 -

marr w or autologous bone marrow, ther was a median of 89-fold expansion of cells (range 18- to 173-fold) after 9 to 19 days of culture. Mean cytotoxicity directed at Daudi and K562 mediated by BMMNC stimulated with OKT3 was 5.5 LU (range 2 - 11) and 4.3 LU (range 2 - 8), respectively. All cultures were
 5 tested 14 days after activation with OKT3 and expanded in the presence of 50 IU/ml of IL-2. Either soluble OKT3 10ng/ml(s) or immobilized OKT3 (coated with 2 µg/ml)(l) were added as indicated.

TABLE 12

10	Source	OKT3 (S or I)	Fold Increase	Daudi	K562
	PBL			LU	LU
	1	I	47	6	3
	2	I	4	11	11
	3	S	12	9	6
	4	S	176	5	4
15	BMMNC				
	1	I	89	2	2
	2	I	18	11	8
	3	S	22	6	5
20	4	S	173	3	2

Anti-CD3 plus anti-CD28 treatment of BMMNC as a source of effector T cells. Proliferative and cytotoxic responses from 3 patients were tested. The patients had received extensive chemotherapy and yet their PBL or BMMNC maintained strong proliferative and cytotoxic responses after anti-CD3 plus anti-
 25 CD28 treatment. PBL or BMMNC (1.5×10^5) were cultured in RPMI plus 5% human serum in the presence of immobilized OKT3 mAb or 50 IU/ml rIL-2 or 9.3 mAb 0.5 µg/ml. Proliferation was assessed on day 3 of culture and cytotoxicity on day 7. Table 13 summarizes the results for BMMNC.

- 63 -

TABLE 13

	SAMPLE	Proliferation ³ H incorp. (cpm)	Cytotoxicity Daudi (LU)	Cytotoxicity K562 (LU)
	Bone Marrow #171 AML, Relapsed			
5	OKT3	23,479	—	—
	OKT3 + anti-CD28 mAb	58,563	18.0	4.8
	OKT3 + IL-2	48,745	15.1	4.6
10	PBL P#632 non-Hodgkins lymphoma, pre-transplant			
	OKT3	58,670	—	—
	OKT3 + anti-CD28 mAb	77,046	14.8	12.7
	OKT3 + IL-2	63,603	18.8	18.1
15	PBL P#635 Hodgkins lymphoma, pre-transplant			
	OKT3	27,758	—	—
	OKT3 + anti-CD28 mAb	46,133	—	—
20	OKT3 + IL-2	43,918	—	—

OKT3-activated T cells (CTC) do not inhibit hematopoietic progenitor growth. In order to determine whether BMMNC mixed with OKT3-activated T cells (CTC) in hematopoietic progenitor assays would inhibit the development of

25 CFU-GM, CTC obtained from PBL after a week of growth was mixed with fresh BMMNC and plated the mixtures into the CFU-GM assay. The autologous CTC were mixed with BMMNC in various ratios, incubated for 1 h at 37°, and then plated in a standard CFU-GM assay. The CTC had no deleterious effect on colony formation, as the number of CFU-GM colonies was within 75% of control

30 over a wide variety CTC:BMMNC ratios (ratios of 1:25 to 5). The number of CFU-GM colonies was not inhibited greater than 90% (an accepted % inhibition of CFU-GM in purged autologous marrow grafts) even at a ratio of 1 CTC to 1

- 64 -

BMMNC. These data suggest that CTC will not inhibit or delay engraftment of autologous bone marrow transplants.

Expansion and cytotoxic functions of PBL or BMMNC from patients before BMT. In order to determine whether PBL or BMMNC from patients heavily pretreated for AML or lymphoma could be activated with anti-CD3 and grown in low dose IL-2, PBL or BMMNC obtained prior to BMT on several patients was tested. The PBL and BMMNC of the patients tested proliferated and exhibited cytotoxicity in a fashion comparable to that seen in normal PBL or BMMNC obtained from normal allogeneic marrow donors. It was anticipated that some patients that had been heavily treated with chemotherapy or radiation would have low counts or have poor responses to anti-CD3 activation. Thus, the number of starting cells was increased in the protocol to compensate for cell loss or inability to proliferate. The use of 9.3 as a costimulant to anti-CD3 activated T cells to enhance helper activity or enhance cytotoxicity could result in improved *in vitro* expansion of activated T cells. Data presented in this example (Specific Example XIV) show that mAb 9.3 can correct proliferative defects in post-transplant lymphocytes further supporting the rationale for using OKT3/9.3 costimulation approach to accelerate immune reconstitution and enhance cytotoxicity directed at malignant cells in ABMT recipients. A recent study by Katsanis, E. *et al.*, *Blood* 78:1286-1291 (1991) shows that T cells from BMT recipients can be expanded by stimulation with OKT3 and IL-2.

Messenger RNA levels for IL-2 receptors (IL-2R), IL-2, and IL-3 in PBL from short and long-term BMT recipients. Earlier studies (see Lum, L.G. *et al.*, *Blood Suppl.* (Abstract) (1991)) suggested that T cells from BMT recipients fail to secrete IL-2 or express IL-2R. Such defects may be due to failure of mRNA synthesis for lymphokines or lymphokine receptors. A determination was made whether T cells from BMT patients failed to express detectable levels of mRNA for IL-2, IL-2R and IL-3. PBL from 11 allogeneic (3 short-term, ST, and 8 long-term, LT) and 4 autologous recipients (2 ST and 2 LT) were tested for levels of IL-2R, IL-2, and IL-3 mRNAs without stimulation (-) or after phytohemagglutinin (PHA) and phorbol ester (TPA) stimulation (+). cDNA synthesized by reverse transcriptase (RTase) from total RNA was amplified by PCR using specific primers and the PCR products run on 1.5% agarose gels containing ethidium

- 65 -

bromide. Table 14 shows the fraction and percent of recipients whose PBL had detectable levels of mRNA for IL-2R, IL-2, and IL-3.

TABLE 14

	STIMULATION	IL-2R (%)	IL-2 (%)	IL-3 (%)
5	Allogeneic Recipients			
	-	2/11 (18)	1/9 (11)	8/11 (73)
	+	9/11 (82)	8/9 (89)	7/10 (70)
	Autologous Recipients			
10	-	2/4 (50)	3/4 (75)	4/4 (100)
	+	4/4 (100)	3/4 (75)	3/4 (75)

PBL from a high proportion of ST and LT autologous and allogeneic BMT recipients expressed levels of mRNAs for IL-2R, IL-2, and IL-3 after stimulation with PHA + TPA. In the ST recipients tested, 2 of 2 ABMT recipients and 3 of 3 allogeneic recipients tested had PBL that expressed mRNA levels of IL-2R and IL-3; 2 of 2 allogeneic recipients tested expressed mRNA for IL-2. In most cases, defective mRNA synthesis for IL-2R, IL-2, and IL-3 may not be responsible for defects in IL-2 secretion and IL-2R expression. Posttranscriptional events may play a more important role in defective lymphokine secretion by T cells from BMT recipients.

CTC help Ig synthesis and express mRNA for lymphokines and perforin.

As discussed in Ueda, M. *et al.*, *J. Cell. Biochem.* (Abstract) (submitted 1992), helper activity was assessed by adding normal T cells or T cells activated with OKT3 to normal B cells after PW stimulation as measured by an ELISA-Plaque (PFC) assay. The number of PFC per million B cells cultured was 3200, 4100, 8800 when 25, 50 or 75 X 10³ normal T cells were added. When the same numbers of CTC were added, the number of PFC were 220, 2100, and 2600. Thus, CTC exhibit substantial helper activity. Furthermore, CTC did not suppress normal autologous or allogeneic T and B cells in a suppressor assay for Ig synthesis. Helper activity was radioresistant. Messenger RNA for IL-2, IL-3, IL-6 and perforin was detected from 6 h to more than 3 days after OKT3 activation using a Rtaase-PCR method. In summary, CTC help B cells produced

- 66 -

Ig and did not suppress Ig synthesis by normal T and B cells. Thus, adoptive transfer of CTC after BMT may not only mediate a GVL effect but may accelerate immune reconstitution.

Defects in anti-CD3-induced proliferative responses in PBL from BMT

- 5 ***patients repaired by adding anti-CD28.*** *In vitro* data on anti-CD3 and anti-CD3/anti-CD28 stimulated proliferative responses of T cells from BMT recipients support the premise that using Mab 9.3 in combination with OKT3 may have potent *in vivo* clinical effects as reported. See Joshi, I. *et al.*, *Blood Suppl.* (Abstract) (1991). T cells from BMT recipients have defects in proliferation after
- 10 mitogen or anti-CD3 stimulation. Previous studies show that costimulation of normal T cells with anti-CD3 (G19-4) and 9.3 enhance anti-CD3-induced proliferation by stabilizing lymphokine mRNAs. Experimentation to assess the ability of anti-CD3 (G19-4 or OKT3) + 9.3 to correct defective anti-CD3-induced proliferative responses in PBL from autologous and allogeneic BMT recipients
- 15 (53-605 days post BMT) was performed. PBL from recipients or controls were stimulated for 3 days with G19-4, G19-4+9.3, OKT3, or OKT3+9.3. 9.3 was added at a final concentration of 100 ng/ml. Fifteen tests were performed on ABMT recipients and sixteen tests were performed on allogeneic recipients. Table 15 shows the number of recipients whose PBL increased (↑), or decreased
- 20 (↓), or did not change (↔) their proliferative responses after the addition of 9.3 to anti-CD3 stimulated PBL. The parenthesis indicate percent of recipients whose proliferative responses increased after the addition of 9.3.

TABLE 15

	BTM RECIPIENTS	CHANGE	TREATMENT	
			G19-4+9.3	OKT3+9.3
25	Autologous BMT	↑	9 (60%)	9 (82%)
		↓	3 (20%)	2 (18%)
		↔	3 (20%)	0 (0%)
30	Allogeneic BMT	↑	11 (69%)	8 (62%)
		↓	5 (31%)	5 (38%)
		↔	0 (0%)	0 (0%)

- 67 -

Costimulation of G19-4+9.3 or OKT3+9.3 significantly increased proliferative responses induced by G19-4 or OKT3 alone ($p < 0.05$, paired rank-sum) in ABMT recipients. In summary, defects in anti-CD3-induced T cell proliferation in BMT recipients were repaired by costimulation with 9.3. These findings have therapeutic implications for patients with immune defects manifest with impaired T cell proliferation. Indeed similar results have been obtained which indicate that the proliferative defect of T cells from patients with HIV infection can be repaired with anti-CD28 treatment. See Lane, H.C. *et al.*, *J. Engl. J. Med.*, 313:85 (1985) regarding proliferate defects in HIV.

10 ***Costimulation with anti-CD3 OKT3 and anti-CD28 9.3 enhanced detectable mRNA levels for IL-2 in PBL from ABMT recipient.*** PBL from a short-term ABMT recipient were studied for expression of mRNA levels for IL-2 after stimulation with OKT3 and costimulation with OKT3/9.3 using RTase-PCR. cDNA synthesized by RTase from total RNA was amplified by PCR using specific
15 primers for IL-2 and the PCR products run on 1.5% agarose gels containing ethidium bromide. Consistent with the findings in the previous paragraph, activated T cells from the ABMT recipient did not have detectable levels of mRNA for IL-2 after OKT3 stimulation alone, whereas the same T cells costimulated with OKT3/9.3 had a distinct band for IL-2 of 458 bp detected on
20 ethidium bromide stained agarose gel. This is an example of how OKT3/9.3 costimulation can repair an apparent defect in the expression of mRNA for IL-2 in T cells from ABMT recipients.

Stimulation of negatively selected CD4⁺ cells with anti-CD28 9.3 after activation with anti-CD3 induces IL-2 independent proliferative responses. CD4⁺
25 cells were purified by a series of negative selection steps as previously described in Thompson, C.B. *et al.*, *PNAS (USA)* 86:1333-1337 (1989). PBL were incubated with a cocktail of mAbs directed at non-CD28⁺ cells, washed, and incubated with immunomagnetic bead coated with goat anti-mouse antibody. The CD28⁺ enriched cells were further purified by removing the CD8⁺ cells by
30 treatment with anti-CD8 and binding the CD8⁺ cells to the immunomagnetic beads.

The remaining CD28⁺, CD4⁺ T cells from a normal donor were cultured by adding cells to culture dishes containing plastic adsorbed OKT3. After 48 h, the cells were removed and placed in flasks containing either rIL-2 (200 IU/ml) or

- 68 -

anti-CD28 mAb (100 ng/ml). The cells were fed with fresh medium as required to maintain a cell density of 0.5×10^6 /ml, and restimulated at approximately weekly intervals by culture on plastic adsorbed OKT3 for 24 h. The cells could be maintained in logarithmic growth, with a 4 to 5 \log_{10} expansion in cells number. As shown in Figure 18, cells propagated with anti-CD3 and anti-CD28 routinely expanded 10 to 30-fold more than cells grown in optimal amounts of anti-CD3 and IL-2. When synthetic medium (X-Vivo 10) not containing FBS was used, anti-CD3 plus anti-CD28 treated cells also expanded 10-fold better than anti-CD3 plus IL-2 treated cells. The highly enriched CD4 cells did not proliferate in the presence of optimal amounts of the lectin phytohemagglutinin (PHA). Thus, the *in vitro* expansion of CD4 cells using anti-CD28 has an advantage over previously described methods, in that it is independent of the addition of exogenous growth factors, as no IL-2 or any other growth factors were added to these cells. In addition, this system does not require the presence of accessory cells, which is advantageous in clinical situations where accessory cells are limiting or defective.

Phenotypes of anti-CD3 activated T cells. Populations of CTC cells grown in IL-2 for 6 to 12 days contained predominantly CD3⁺ cells (greater than 84%, median 88%). The proportion of CD56⁺ cells (a marker for NK cells) was less than 1.3%. Triggering of E⁺ cells with OKT3 is preferentially selecting CD3⁺ cells. CD4⁺ cells were 18% or less and CD8⁺ cells were greater than 66%. Lytic activity did not correlate with the proportions of CD56⁺ cells in the cultures.

Immunophenotype of T cells differs after anti-CD28 and IL-2-mediated cellular growth. To examine the subsets of T cells that are expanded, PBL were propagated for 16 days using either anti-CD3 and IL-2 or anti-CD3 and anti-CD28. The percentage of CD4 and CD8 cells was 23.8 and 84.5 in the cells grown in IL-2, and 56.0 and 52.6 in the cells grown in CD28. These results suggests that CD28 expansion favors the CD4⁺ cells, in contrast to the well established observation that CD8⁺ cells predominate in cells grown in IL-2 (for example, see above paragraph; see also Cantrell, D.A. *et al.*, *J. Exp. Med.* 158:1895 (1983)). To further test this possibility, CD4 cells were enriched to 98% purity using negative selection with monoclonal antibodies and magnetic immunobeads as described elsewhere in this example. The cells were cultured

- 69 -

for one month using anti-CD3 and either IL-2 or anti-CD28 to propagate the cells. There was equal expansion of the cells for the first 26 days of the culture, however, as can be seen in Table 16, the phenotype of cells diverged progressively with increasing time in culture.

5

TABLE 16

	CULTURE METHOD	DAYS	CD3 (%)	CD4 (%)	CD8 (%)
10	anti-CD3 + IL-2	0	>99	>99	<1
		6	>99	98	1
		12	>99	85	10
		20	>99	45	40
		26	>99	12	78
15	anti-CD3 + IL-2	0	>99	>99	<1
		6	>99	>99	<1
		12	>99	>99	<1
		20	>99	>99	<1
		26	>99	>99	<1

Use of anti-CD28 for in vitro expansion of TIL. The use of IL-2 and inactivated tumor cells to expand tumor infiltrating lymphocytes (TIL cells) for later adoptive immunotherapy with a variety of neoplasms has demonstrated promise (*e.g.*, see Rosenberg, S.A. *et al.*, *NEJM* 323:570-578 (1990)). TIL cells were isolated from a nephrectomy specimen from a patient with renal cell carcinoma. The cells were cultured with tumor cells, and either IL-2 or anti-CD28 and IL-2 with mAb OKT3 added at weekly intervals, beginning at day 14. Table 17 demonstrates that anti-CD28 is an improved method for the propagation of these cells in some patients, with a 20-fold greater yield of cells. Immunophenotype analysis also reveals that CD4 T cells are expanded in "TIL" cultures. Furthermore, these cells also exhibited potent cytotoxic activity against DAUDI targets, with 82.8, 69.7, 78.8 and 101.5 percent specific lysis at effector to target ratios of 40:1, 20:1, 10:1 and 5:1.

- 70 -

TABLE 17

Use of anti-CD28 to Expand TIL cells

Day of Culture		Tumor cells + IL-2 (1000 U/ml)	Tumor cells + IL-2 anti-CD3, then anti- CD28
5	0	3.6×10^7	3.6×10^7
	5	5.8×10^7	1.3×10^8
	12	1.7×10^8	1.4×10^8
	16	2.0×10^8	2.6×10^8
	19	3.8×10^8	4.8×10^8
10	25	2.9×10^8	9.0×10^8
	36	3.7×10^8	1.8×10^9
	43	3.0×10^8	3.2×10^9
	48	2.3×10^8	5.1×10^9

Anti-CD3 and anti-CD28 costimulation enhances expression of mRNA for

- 15 ***IL-2 and TNF- α in CD4⁺ cells.*** To examine whether CD4 cells propagated *in vitro* by anti-CD28 might be an effective source of lymphokines, resting CD4 cells were stimulated by anti-CD3 mAb for 48 h followed by the addition of 50 IU/ml of recombinant IL-2 and compared with CD4 T cells costimulated with anti-CD3 and anti-CD28. Total RNA was harvested from each combination. A
- 20 total of 10 μ g of RNA was loaded into each lane. A class I probe (HLA B7) was used to show uniform loading. The blot was hybridized with ³²P labeled probes specific for IL-2, TNF- α and HLA in succession. On days 1 and 8, the cultures were restimulated with anti-CD3 mAb in the presence of IL-2 or anti-CD28 mAb 9.3 (0.1 μ g/ml). The blots were stripped, and rehybridized with a probe for a
- 25 constant region of HLA class I mRNA, to demonstrate equal loading of the lanes. As illustrated by the Northern Blot of Figure 19, there was a clear enhancement of mRNA for IL-2 and TNF- α after costimulation with anti-CD3 and anti-CD28 and the IL-2 and TNF- α mRNA exceeded that of the IL-2 propagated cells by 10 to 50-fold. Similar results were obtained when the culture was examined for
- 30 one month of culture, after weekly restimulation with anti-CD3 and IL-2 or anti-CD28.

- 71 -

Supernatants from long-term anti-CD3 and anti-CD28 stimulated cultures of CD4⁺ cells contain substantial amounts of IL-2, GM-CSF, and TNF- α . After anti-CD3 stimulation in the presence of 200 IU/ml of IL-2 or the combination of anti-CD3 and anti-CD28 in the absence of IL-2, supernatants were tested for IL-2 content using the CTLL-2 cell line, GM-CSF content by ELISA, and TNF- α content by ELISA. The CD4⁺ cells were restimulated with anti-CD3 and IL-2 or anti-CD3 and anti-CD28 respectively at approximately weekly intervals. Anti-CD3 and anti-CD28 cultures of CD4⁺ cells produced roughly the same amount of IL-2 found in supernatant obtained from anti-CD3 activated cells grown in exogenous IL-2. The amount of GM-CSF produced by anti-CD3 and anti-CD28 stimulated CD4⁺ cells was also substantial. Although there were variations in levels of TNF- α depending on when the supernatants were tested, costimulation with anti-CD3 and anti-CD28 was superior to stimulation with anti-CD3 and IL-2 for inducing mRNA for TNF- α . These data indicate that anti-CD28 costimulation with anti-CD3 may not only replace some of the functions of IL-2 but may enhance other synthetic functions of CD4⁺ cells.

SPECIFIC EXAMPLE XV

CD28 and CTLA-4 Expression Studies.

CTLA-4 expression limited to CD28⁺ T cells. Utilizing site-specific primers and DNA PCR of human genomic DNA, a 348 bp fragment corresponding to exon II of human CTLA-4 was generated, gel purified and used as a ³²P-labeled probe. Purified human T cells were separated into CD28⁺ and CD28⁻ fractions by negative selection with magnetic bead immunoabsorption. CD28⁻ T cells were either tested in media or stimulated with PMA + ionomycin or PMA + anti-CD28 mAb (mAb 9.3) for 12 h. CD28⁺ cells were stimulated with the last two conditions for 12 h. RNA was extracted by guanidinium isothiocyanate and purified over cesium chloride gradients. Equal amounts of RNA (as determined by ethidium bromide staining) were loaded and separated on a formaldehyde-agarose gel and transferred to nitrocellulose to demonstrate equal loading of RNA. This blot was subsequently probed with the CTLA-4 probe generated above. CTLA-4 was expressed in CD28⁺ cells following PMA or PMA + mAb 9.3 stimulation but not expressed in resting or stimulated CD28⁻ cells. The same blot was hybridized to a HLA probe to confirm equal loading of RNA.

- 72 -

The expression of CTLA-4 induced under conditions causing CD28

pathway activation. Purified resting CD28⁺ T cells were stimulated with PMA alone, ionomycin alone, and PMA + ionomycin for 1 h, 6 h, 12 h and 24 h. RNA was extracted and analyzed by hybridization to CD28 and CTLA-4 probes.

- 5 Northern blot analysis showed that CTLA-4 expression was induced by PMA or PMA + ionomycin, conditions which are costimulatory with CD28 pathway activation. CTLA-4 expression was not induced by ionomycin, which is not costimulatory with CD28 pathway activation. In contrast, CD28 expression was constant with ionomycin or PMA and even appeared to be suppressed with PMA
10 and ionomycin stimulation. It should also be noted that the induction of CTLA-4 expression occurred as soon as 1 h after stimulation, compared to 6 - 12 h with IL-2 expression following CD28 pathway activation. Since expression of CTLA-4 precedes the biological events caused by CD28 pathway activation (i.e. enhanced IL-2 expression), CTLA-4 expression likely plays a role in the
15 generation of later events.

- Purified human T cells were either untreated or stimulated for 1 h, 4 h or 23 h with PMA + PHA, anti-CD28 mAb crosslinked with a second antibody (goat anti-mouse Ig), or anti-CD5 mAb crosslinked in the same manner. Northern blot analysis showed that crosslinking of CD28 receptors, which also
20 can activate the CD28 pathway through mechanisms distinct from PMA and ionomycin, also induced CTLA-4 expression.

- CD28⁺ cell lines slightly or not responsive to CD28 pathway activation do not express CTLA-4.*** Two cell lines that are CD28⁺ but responded poorly (T cell line Jurkat CJ) or not at all (myeloma cell line RPMI-8226) to CD28
25 costimulation as discussed in Kozbor, D. *et al.*, *J. Immunol.*, 138:4128-4132 (1987) and Ledbetter, J.A. *et al.*, *PNAS (USA)*, 84:1384-1388 (1987), were stimulated with PMA + ionomycin + mAb 9.3 and subsequently analyzed by Northern blot for CD28 and CTLA-4 expression. Northern blot analysis of the T cell leukemia cell line Jurkat CJ and of the myeloma cell line RPMI 8226 showed
30 that these cell lines did not express CTLA-4 despite CD28 expression.

T cells were incubated with various combinations of mitogens including phytohemagglutinin (PHA), phorbol myristate acetate (PMA), and ionomycin (IONO), or anti-CD28 monoclonal antibodies (α -CD28), and examined for the ability to induce CTLA-4 mRNA expression. As shown in Figure 20, the

- 73 -

combination of the mitogens PHA and PMA readily induce CTLA-4 expression in normal human T cells, but all combinations tested failed to induce the expression of the CD28-isoform CTLA-4 in the Jurkat T cell line. Both cell types express significant levels of CD28 mRNA under all conditions tested.

- 5 Resting T cells were costimulated with anti-CD28 monoclonal antibodies to examine activation in normal human cells. As shown in Figure 21, normal human T cells stimulated by plate-adherent anti-CD3 (α -CD3) monoclonal antibodies at a concentration of 1 μ g/ml induced only low levels of CTLA-4 mRNA expression first observable at 1 h after stimulation. In contrast,
- 10 costimulation of cells with anti-CD3 monoclonal antibodies (1 μ g/ml) soluble anti-CD28 (α -CD28) monoclonal antibody (1 μ g/ml) led to a dramatic increase in the induction of CTLA-4 mRNA expression.

SPECIFIC EXAMPLE XVI

Efficacy of Administration of CTLA-4Ig as a Treatment for Autoimmune Disease.

- 15 Experimental Autoimmune Encephalomyelitis (EAE) is a rodent and primate model for multiple sclerosis. Data has been generated on the effect of administration of CTLA-4Ig in both passive (indirect) and active (direct) models of EAE. CTLA-4Ig is a fusion protein consisting of the extracellular domain of human CTLA-4 fused to the constant region of human IgG1 (referred to here as
- 20 huCTLA-4Ig).

- Adoptively Transferred (Passive) EAE.* In the passive EAE model, donor mice are immunized with 0.4 mg Myelin Basic Protein (MBP) in Complete Freund's Adjuvant (CFA), divided over four quadrants. The draining axillary and inguinal lymph nodes are removed eleven days later. Lymph node cells (4 x
- 25 10^6 /ml) are plated in 2 ml cultures in 24 well plates, in the presence of 25 μ g/ml MBP. After four days in culture, 30×10^6 of the treated cells are injected into the tail vein of each naive, syngeneic recipient mouse.

The recipient mice develop a remitting, relapsing disease and are evaluated utilizing the following criteria:

- 74 -

- 0 - normal, healthy
1 - limp tail, incontinence; occasionally the first sign of the disease is a "tilt"
2 - hind limb weakness, clumsiness
3 - mild paraparesis
4 - severe paraparesis
5 - quadriplegia
6 - death

Using this passive model of EAE, the effect of huCTLA-4Ig treatment of the donor cells on disease severity was tested in PLSJLF1/J mice. Treatment of lymph node cells *in vitro* with MBP was performed either in the presence or the absence of 30 μ g/ml huCTLA-4Ig. The treated cells were then introduced into a syngeneic recipient mouse. As shown in Figure 22, mice receiving huCTLA-4Ig-treated cells (designated PPIB CTLA-4) showed a significantly reduced severity of their first episode of disease as compared to mice receiving untreated cells (designated PPIA control). In addition, ensuing relapses in the mice receiving huCTLA-4Ig-treated cells were less severe than in mice receiving cells not exposed to huCTLA-4Ig. In fact, all five mice receiving huCTLA-4Ig-treated cells stopped relapsing, and no longer showed signs of disease at 80-100 days post transfer.

Clinical disease severity was reduced even further by treating both the donor mice and the cultured cells with huCTLA-4Ig (Figure 23). In these experiments, donor mice of the SJL/J strain were given either 100 μ g huCTLA-4Ig or 100 μ g chimeric control IgG1 intraperitoneally each day for eleven days. T cells were then isolated from lymph nodes of these donors and cultured with MBP *in vitro* in the presence of either 30 μ g/ml huCTLA-4Ig or chimeric control IgG1. The treated cells were then introduced into a syngeneic recipient. Treatment of either the donor mice or the *in vitro* cultures resulted in significantly reduced clinical disease severity. Treatment of both the donor mice and the cultured cells with huCTLA-4Ig was the most effective protocol for reducing clinical disease severity.

Direct administration of huCTLA-4Ig to mice receiving adoptively transferred cells was also examined. As shown in Figure 24, when PLSJLF1/J recipient mice were given 100 μ g of either huCTLA-4Ig or human IgG in PBS intraperitoneally on days 1 to 9 post transfer, no difference in disease severity was observed between the two groups of mice. However, in experiments

- 75 -

utilizing SJL/J mice, reduced disease severity during relapse was noted in mice treated with 100 μ g huCTLA-4Ig intraperitoneally on days 1 to 5 post transfer (Figure 25). Ongoing experiments are examining the effect of administration of a single dose of huCTLA-4Ig to SJL/J recipient mice on day 2 post transfer.

- 5 Table 18 shows results of such an experiment, compared to the results obtained when recipient mice are given either huCTLA-4Ig or human IgG1 on days 1 to 5 post transfer. While severity of the first episode of disease did not appear to be significantly altered by treatment with huCTLA-4Ig either on day 2 or days 1-5, the duration of the first episode of disease was shorter for mice
10 given huCTLA-4Ig treatment (Table 18). In addition, huCTLA-4Ig treatment on days 1-5 resulted in delayed onset of the first episode of disease.

TABLE 18

15 Effect of Administration of CTLA-4Ig to Mice with Adoptively Transferred EAE

	CTLA-4Ig (day 2)	CTLA-4Ig (days 1-5)	IgG Control (days 1-5)
Mean day of onset	11.3	16.3	10.8
Maximum score of first episode	3.2	2.8	2.8
20 Maximum score to date (day 35)	3.5	3.2	3.0
Average of first episode (days)	5.2	6.7	9.8

- 25 **Direct (Active) Model of EAE.** Studies using a direct (active) model of EAE have also been conducted. In these experiments, huCTLA-4Ig was directly administered to mice immunized with MBP and treated with pertussis toxin (PT). PLSJLFI/J mice immunized with MBP on day 0 and injected with PT
30 intravenously on days 0 and 2 were given either huCTLA-4Ig or control IgG1 on days 0 to 7. Five of ten mice given huCTLA-4Ig died on days 5 to 7, for reasons related to the PT administered, and not the experimental design. The results of the experiment for the remaining five mice are as shown in Figure 27. Administration of huCTLA-4Ig markedly reduced the mean clinical severity of
35 disease in these animals, as compared to the mice treated with IgG1. These

- 76 -

findings indicate that direct administration of soluble human huCTLA-4Ig can provide an effective therapeutic strategy in the treatment of autoimmune disease.

5 It should be appreciated that a latitude of modification, change or substitution is intended in the foregoing disclosure and, accordingly, it is appropriate that the appended claims be construed broadly and in a manner consistent with the spirit and scope of the invention herein.

All publications and applications cited herein are incorporated by reference.

- 77 -

WHAT IS CLAIMED IS:

1. A method of inhibiting CD28 pathway activation associated with an increase in cellular production of a T_H CD28 lymphokine in a CD28⁺ T cell population, such activation occurring upon binding of a stimulatory CD28 ligand to a CD28 receptor stimulatory binding site on T cells in the population, the method comprising the steps of:
 - a) selecting an inhibitory ligand capable of binding to the stimulatory CD28 ligand;
 - b) providing the inhibitory ligand in a biologically compatible form; and
 - c) administering the inhibitory ligand to the population in an amount sufficient to bind and inhibit the stimulatory ligand from binding the CD28 receptor stimulatory binding site.
2. The method of Claim 1, wherein the stimulatory ligand comprises a natural CD28 ligand.
3. The method of Claim 1, wherein the inhibitory ligand comprises an antibody to the stimulatory ligand or a fragment of an antibody to the stimulatory ligand.
4. The method of Claim 1, wherein the inhibitory ligand comprises a soluble form of CTLA-4.
5. The method of Claim 4, wherein the inhibitory ligand comprises CTLA-4Ig.
6. The method of Claim 1, wherein the inhibitory ligand is of synthetic origin.
7. The method of Claim 1, wherein the inhibitory ligand comprises a recombinant molecule.

- 78 -

8. The method of Claim 1, further comprising the step of:
- d) administering to the population a second inhibitory ligand capable of binding but not stimulating the CD28 receptor binding site.

9. A method of suppressing the production of a T_H CD28 lymphokine by a population of CD28⁺ T cells, the method comprising the steps of:

- a) providing an inhibitory ligand which binds a stimulatory ligand for CD28;
- b) providing the ligand in biologically compatible form; and
- c) administering the provided ligand in an amount sufficient to suppress production of the lymphokine in the population.

10. The method of Claim 9, wherein the inhibitory ligand comprises a soluble form of CTLA-4.

11. The method of Claim 10, wherein the inhibitory ligand comprises CTLA-4Ig.

12. The method of Claim 9, wherein the T cell population is in a patient in an autoimmune state.

13. A method of suppressing T_H CD28 lymphokine production in a patient having a population of CD28⁺ T cells, the method comprising the steps of:

- a) providing an inhibitory ligand which binds a natural stimulatory ligand for CD28; and
- b) administering the inhibitory ligand in a therapeutically effective amount to the population of T cells.

14. The method of Claim 13, wherein the step of administering the inhibitory ligand to the population of T cells is carried out *in vivo*.

- 79 -

15. The method of Claim 13, wherein the step of administering the ligand to the population of T cells is carried out *in vitro*, and further comprising the step of:

d) after the administering step, introducing the population of T cells into the patient.

16. The method of Claim 15, wherein the T cell population is removed from the patient prior to administering the inhibitory ligand to the T cell population.

17. The method of Claim 13, wherein the inhibitory ligand comprises a soluble form of CTLA-4.

18. The method of Claim 17, wherein the inhibitory ligand comprises CTLA-4Ig.

19. A method of treating an autoimmune disease in a patient comprising the steps of:

a) selecting an inhibitory ligand which binds a natural stimulatory ligand to CD28; and

b) administering a therapeutically effective amount of the inhibitory ligand to the patient.

20. The method of Claim 19, wherein the inhibitory ligand comprises an antibody to the stimulatory ligand or a fragment of an antibody to the stimulatory ligand.

21. The method of claim 19, wherein the stimulatory ligand is of synthetic origin.

22. The method of claim 19, wherein the stimulatory ligand comprises a recombinant molecule.

- 80 -

23. The method of Claim 19, further comprising the step of:

- d) administering to the patient a second inhibitory ligand capable of binding but not stimulating the CD28 receptor binding site.

24. The method of Claim 19, wherein the stimulatory ligand is B7 and the inhibitory ligand comprises a soluble form of CTLA-4.

25. The method of Claim 19, wherein the inhibitory ligand comprises CTLA-4lg.

26. The method of Claim 19, wherein the step of administering the ligand to the patient is carried out by administering the ligand to a population of T cells *in vitro*, and further comprises introducing the population of T cells into the patient.

27. The method of Claim 26, wherein the T cell population is removed from the patient prior to administering the inhibitory ligand to the T cell population.

28. The method of Claim 19, wherein the step of administering the inhibitory ligand to the patient is carried out *in vivo*.

29. The method of Claim 19, wherein the step of administering the inhibitory ligand to the patient is carried out *in vitro* by administering the inhibitory ligand to a population of cells removed from the patient, and further comprising the step of:

- c) thereafter reintroducing the cells to the patient.

30. The method of Claim 24, wherein the autoimmune disease is multiple sclerosis.

31. The method of claim 28, whereby the administration of the inhibitory ligand is by injection.

- 81 -

32. The method of claim 29, wherein the population of cells comprises lymph node cells.

33. The use in manufacture of a medicament for treating an autoimmune disease of an inhibitory ligand capable of binding but not stimulating the extracellular CD28 receptor binding site.

1/22




FIG. 1




FIG. 2

SUBSTITUTE SHEET (RULE 26)




FIG. 3





FIG. 5

**FIG. 4**

SUBSTITUTE SHEET (RULE 26)

4 / 22




FIG. 6A




FIG. 6B

SUBSTITUTE SHEET (RULE 26)

5/22




FIG. 7A






FIG. 7B

6/22

**FIG. 8A****FIG. 8B**

7/22




FIG. 9A




FIG. 9B

8/22




FIG. 10A




FIG. 10B

9/22




FIG. 11




FIG. 13

10/22




FIG. 12A




FIG. 12B

11/22




FIG. 14




FIG. 15

12 / 22




FIG. 16




FIG. 17

13/22




FIG. 18




FIG. 19




FIG. 20
SUBSTITUTE SHEET (RULE 26)






FIG. 21

17/ 22

**FIG. 22**

18 / 22

**FIG. 23**

SUBSTITUTE SHEET (RULE 26)

19/ 22




FIG. 24

20/22




FIG. 25

21/22




FIG. 26

22 / 22




FIG. 27

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/06701

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 35/14, 37/00, 39/00; C07K 3/00, 13/00, 15/00, 17/00; C12P 21/06

US CL : 424/130.1; 435/69.7; 514/12; 530/350, 387.3, 388.33, 388.75, 395

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1; 435/69.7; 514/12; 530/350, 387.3, 388.33, 388.75, 395

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. IMMUNOLOGY, VOLUME 138, NUMBER 12, ISSUED 15 JUNE 1987, KOZBOR ET AL., "Tp44 MOLECULES INVOLVED IN ANTIGEN-INDEPENDENT T CELL ACTIVATION ARE EXPRESSED ON HUMAN PLASMA CELLS", PAGES 4128-4132, SEE ENTIRE DOCUMENT	1-33
Y	J. IMMUNOLOGY, VOLUME 136, NUMBER 9, ISSUED 01 MAY 1986, MARTIN ET AL., "A 44 KILODALTON CELL SURFACE HOMODIMER REGULATES INTERLEUKIN 2 PRODUCTION BY ACTIVATED HUMAN T LYMPHOCYTES", PAGES 3282-3287, SEE ENTIRE DOCUMENT.	1-33

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

"	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance		
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

19 JULY 1994

Date of mailing of the international search report

AUG 01 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

DONALD E. ADAMS, PH.D.

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06701

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. IMMUNOLOGY, VOLUME 135, NUMBER 4, ISSUED OCTOBER 1985, LEDBETTER ET AL., "ANTIBODIES TO Tp67 AND Tp44 AUGMENT AND SUSTAIN PROLIFERATIVE RESPONSES OF ACTIVATED T CELLS", PAGES 2331-2336, SEE ENTIRE DOCUMENT.	1-33
Y	J. EXP. MED., VOLUME 174, ISSUED SEPTEMBER 1991, LINSLEY, ET AL., "CTLA-4 IS A SECOND RECEPTOR FOR THE B CELL ACTIVATION ANTIGEN B7", PAGES 561-569, SEE ENTIRE DOCUMENT.	1-33
Y	CANCER IMMUNOL. IMMUNOTHER., VOLUME 34, ISSUED 1992, BASSE ET AL., "TISSUE DISTRIBUTION OF ADOPTIVELY TRANSFERRED ADHERENT LYMPHOKINE-ACTIVATED KILLER CELLS ASSESSED BY DIFFERENT CELL LABELS", PAGES 221-227, SEE ENTIRE DOCUMENT.	1-33
Y	IMMUNOLOGY TODAY, VOLUME 11, NUMBER 6, ISSUED 1990, OSBAND ET AL., "PROBLEMS IN THE INVESTIGATIONAL STUDY AND CLINICAL USE OF CANCER IMMUNOTHERAPY", PAGES 193-195, SEE ENTIRE DOCUMENT.	1-33
Y	ANNALS OF INTERNAL MEDICINE, VOLUME 111, NUMBER 7, ISSUED 01 OCTOBER 1989, DILLMAN, "MONOCLONAL ANTIBODIES FOR TREATING CANCER", PAGES 592-600, SEE ENTIRE DOCUMENT.	1-33
Y	CARNEY ET AL., "GENES AND CANCER", PUBLISHED 1990 BY JOHN WILEY & SONS LTD. (N.Y.), CHAPTER 17, PAGES 183-189, SEE ENTIRE DOCUMENT.	1-33
Y	TIBTECH, VOLUME 11, ISSUED FEBRUARY 1993, HARRIS ET AL., "THERAPEUTIC ANTIBODIES - THE COMING OF AGE", PAGES 42-46, SEE ENTIRE DOCUMENT.	1-33
Y	SCIENCE, VOLUME 252, ISSUED 21 JUNE 1991, WALDMANN, "MONOCLONAL ANTIBODIES IN DIAGNOSIS AND THERAPY", PAGES 1657-1662, SEE ENTIRE DOCUMENT.	1-33